

From the Department of Medical Biochemistry and Biophysics
Karolinska Institutet, Stockholm, Sweden

EXPLORING THE ROLE OF PDGF-D IN HEALTH AND DISEASE

Hanna Gladh



**Karolinska
Institutet**

Stockholm 2016

All previously published papers were reproduced with permission from the publisher.

Published by Karolinska Institutet.

Printed by Eprint AB 2016

© Hanna Gladh, 2016

ISBN 978-91-7676-297-4

Exploring the role of PDGF-D in health and disease

THESIS FOR DOCTORAL DEGREE (Ph.D.)

By

Hanna Gladh

Principal Supervisor:

Professor Ulf Eriksson
Karolinska Institutet
Department of Medical Biochemistry and
Biophysics
Division of Vascular biology

Co-supervisor:

Erika Folestad
Karolinska Institutet
Department of Medical Biochemistry and
Biophysics
Division of Vascular biology

Opponent:

Professor Aristidis Moustakas
Uppsala University,
Department of Medical Biochemistry and
Microbiology

Examination Board:

Docent Jorge Ruas
Karolinska Institutet
Department of Physiology and Pharmacology

Docent Johan Kreuger
Uppsala University
Department of Medical Cell Biology

Docent Mats Hellström
Uppsala University
Department of Immunology, Genetics and
Pathology

Till Morfar

POPULÄRVETENSKAPLIG SAMMANFATTNING

Flercelliga organismer är uppbyggda av flera sorters celler med många olika funktioner. Vissa av cellerna är specialiserade på att få hjärtat att slå, medan andra celler står beredda ifall ett sår behöver läkas, eller ser till att blod kan transporteras så att alla organismens celler kan få näring. För att organismens celler ska fungera på ett korrekt sätt behöver cellerna kommunicera med varandra. Detta sker via signalsubstanser som utsöndras av den cell som vill något. Signalen tas emot av en mottagare - en receptor, som finns på mottagarcellen. Mottagarcellen reagerar på signalen, t.ex. genom att dela sig, röra på sig eller skicka ut nya signaler. Ibland kan det bli fel i cellernas kommunikation, vilket är vanligt i olika sjukdomssammanhang. Felet kan bero på att cellerna har blivit skadade så att de inte längre kan signalera normalt, som vid cancer, då enskilda celler förlorar förmågan att svara på vissa signaler, med följden att de börjar växa på ett okontrollerat sätt.

En familj av signalsubstanser är PDGF (Platelet-Derived Growth Factor), som normalt utsöndras från bland annat blodkärlsceller. PDGF-familjen består av fyra olika typer av PDGF substanser som alla har olika funktioner i organismen, men gemensamt är att de fungerar som tillväxtfaktorer. PDGF-signaleringsen är livsviktig under fosterstadiet, där specifika celler får signaler om hur de ska dela sig och vandra till platser där de behövs. Utan korrekt PDGF-signaleringsen utvecklas därför inte organismen som den ska. En annan viktig funktion för PDGF är att se till att blodkärlen är omslutna av stödjande celler, så att de håller ihop och inte läcker. Dessutom är PDGF involverat i cancer, där för mycket PDGF-signaleringsen kan leda till att det bildas cancerceller, eller att cancerceller stimuleras att växa.

Den här avhandlingen handlar om PDGF-D, den senast upptäckta medlemmen i PDGF-familjen, som även är den substans man vet minst om. För att försöka förstå vad PDGF-D gör har vi konstruerat en genetiskt modifierad musstam som saknar PDGF-D. I artikel I undersöker vi den musstammen och visar att PDGF-D inte är nödvändigt för normal fosterutveckling, men att avsaknaden av PDGF-D resulterar i ett lite förhöjt blodtryck. Vi visar även att PDGF-D finns i blodkärl, både i cellagret som utgör själva "röret" och i stödjeceller som sitter runt om, samt att vissa av stödjecellerna är beroende av PDGF-D för att fungera optimalt. I Artikel II visar vi att en speciell form av cancer i bukspottkörteln som saknar PDGF-D växer mindre. Vi kom fram till att det berodde på att PDGF-D signaleringsen påverkar en mycket liten cellpopulation, som påverkar tumörens tillväxt och utveckling. Slutligen, i Artikel III fokuserar vi på NRP1, som fungerar som en tilläggsreceptor till en annan receptor för PDGF-D och vi visar att NRP1 kan påverka PDGF-D signaleringsen.

De forskningsfynd som presenteras i avhandlingen utvidgar kunskapen om hur PDGF-familjen fungerar och signalerar. Den ökade förståelsen för hur cellerna kommunicerar med hjälp av PDGF behövs för att kunna utveckla nya, bättre läkemedel mot sjukdomar där bland annat blodkärlen är påverkade, till exempel åderförfettning och cancer.

ABSTRACT

The platelet-derived growth factors (PDGF) and their receptors (PDGFRs) regulate growth and migration in cell populations of mesenchymal origin. The PDGF signaling system is vital for development, in neural crest formation, in organogenesis, and in maturation of the microvasculature. In the adult, PDGFs are needed to maintain homeostasis. They are also released in response to tissue injury, where they promote wound healing and neovascularization. In the adult, high expression of PDGFs is also seen in atherosclerosis, fibrosis and in malignant conditions. The PDGF family consists of four ligands that are present as dimers (PDGF-AA, PDGF-BB, PDGF-CC and PDGF-DD) and two tyrosine kinase receptors (PDGFR α and PDGFR β). Upon ligand binding, receptor dimerization and auto-phosphorylation is induced. Downstream signaling leads to immediate effects in receptor-expressing cells, but also prolonged effects through modulation of transcription are seen.

PDGF-D is the most recently found ligand and its biological function is still unclear, although its signaling receptor PDGFR β is mainly expressed in vascular smooth muscle cells, thus indicating a vascular role also for PDGF-D. PDGF-B, the other PDGFR β ligand, also binds to PDGFR α , thus making PDGF-D the only ligand that signals exclusively through PDGFR β . Moreover, PDGF-D expression is uncoupled from its signaling as it is released in a latent, full-length form requiring proteolytic cleavage for receptor binding. In contrast, the other PDGFR β ligand, PDGF-B, is active already upon release.

PDGF signaling has been studied through a multitude of genetically modified animals, and these studies have contributed greatly to the understanding of PDGF function. In the work included in this thesis, we present the PDGF-D knockout mouse strain, and characterize the expression and function of PDGF-D *in vivo*, in both physiological conditions and in the tumor setting. We confirm that PDGF-D has a vascular expression pattern, and show that it is mainly expressed in arteries and in the endothelium, but it can also be expressed in vSMCs. We show that targeted deletion of PDGF-D affected an NG2-expressing pericyte population in the heart, and that animals lacking PDGF-D have slightly elevated blood pressure. Furthermore, we present evidence that paracrine PDGF-D signaling from the vasculature induces the production of factors from a rare PDGFR β -expressing tumor cell subpopulation, thereby contributing to tumor growth. We also define a possible role for a co-receptor in this process. Finally, we present NRP1 as co-receptor for PDGF-D in PDGFR β signaling, and thereby also suggest a mechanistic basis for PDGF-D-specific PDGFR β -NRP1 complex formation and signaling. The addition of NRP1-mediated modulation adds complexity to the current model of PDGF-D/PDGFR β signaling. Ultimately, these findings will lead to a better understanding of the role(s) of PDGF-D signaling, and thereby to improved development of tailored therapeutics for conditions where PDGF-D signaling might be dysregulated, such as atherosclerosis and cancer.

LIST OF SCIENTIFIC PAPERS

This thesis is based on the following papers, which will be referred to in the text by their roman numerals:

- I. **Gladh H***, Folestad EB*, Muhl L, Ehnman M, Tannenbergs P, Lawrence AL, Betsholtz C and Eriksson U. Mice lacking Platelet-Derived Growth Factor D display a mild vascular phenotype.
PLoS One. 2016 Mar 31;11(3):e0152276. (*Equal contribution)
- II. Cortez E, **Gladh H**, Braun S, Bocci M, Cordero E, Björkström NK, Miyazaki H, Michael IP, Eriksson U, Folestad E and Pietras K. Functional malignant cell heterogeneity in pancreatic neuroendocrine tumors revealed by targeting of PDGF-DD.
Proc Natl Acad Sci U S A. 2016 Feb 16;113(7):E864-73
- III. Muhl L, **Gladh H**, Folestad EB, Wang Y, Jakobsson L, Eriksson U. Neuropilin 1 is a co-receptor for Platelet-Derived Growth Factor (PDGF)-D/PDGF receptor (PDGFR) β signaling.
Manuscript

CONTENTS

1	Introduction.....	1
1.1	The PDGF system.....	1
1.1.1	PDGF ligands, structure and activation.....	1
1.1.2	PDGF receptors, structure and signaling.....	2
1.2	Functional aspects of PDGF signaling.....	5
1.2.1	Expression and regulation	5
1.2.2	Physiological roles.....	7
1.3	Pathological aspects of PDGF signaling.....	11
1.3.1	Fibrosis.....	11
1.3.2	Atherosclerosis and other vascular pathologies	12
1.3.3	Cancer	13
1.4	Neuropilin 1	17
1.4.1	Structure, function and expression of NRP1	17
1.4.2	NRP1-mediated modulation of signaling.....	18
2	Aims of this thesis.....	21
3	Papers and discussion	23
3.1	Paper I – Mice Lacking Platelet-Derived Growth Factor D Display A Mild Vascular Phenotype	23
3.2	Paper II – Functional malignant cell heterogeneity in pancreatic neuroendocrine tumors revealed by targeting of PDGF-DD	24
3.3	Paper III – Neuropilin 1 is a co-receptor for Platelet-Derived Growth Factor (PDGF)-D/PDGF Receptor (PDGFR) β signaling	26
4	Future perspectives	29
5	Acknowledgements.....	31
6	References.....	35

LIST OF ABBREVIATIONS

CAF	Cancer-associated fibroblast
CS	Chondroitin sulfate
CSC	Cancer stem cell
CUB	<u>C</u> omplement subcomponents c1r/c1s, <u>U</u> rchin EGF-like protein and <u>B</u> one morphogenic protein 1
ECM	Extracellular matrix
EMT	Epithelial-to-mesenchymal transition
FACS	Fluorescence-activated cell sorting
GFD	Growth factor domain
LRP	Low-density lipo-protein receptor related protein
MMP	Matrix metalloproteinases
M _r	Relative molecular weight
mRNA	Messenger ribonucleic acid
NG2	Nerve/glial antigen 2
NRP	Neuropilin
PanNET	Pancreatic neuroendocrine tumors
PDGF	Platelet-derived growth factor
PDGFR	Platelet-derived growth factor receptor
PI3K	Phosphatidylinositol 3'-kinase
SEMA3	Class-3 semaphorin
SH2	Src homology 2
TGF- β	Transforming growth factor β
TIMP	Tissue inhibitor of metalloproteinase
tPA	Tissue plasminogen activator
uPA	Urokinase plasminogen activator
VEGF	Vascular endothelial growth factor
VEGFR	Vascular endothelial growth factor receptor

1 INTRODUCTION

The family of platelet-derived growth factors (PDGFs) is important during development, in adult physiology, and in a number of pathological conditions, including atherosclerosis, fibrosis, and cancer. They act by inducing survival, proliferation and migration in cells of mesenchymal origin, and are often released upon tissue injury (reviewed in (1)). Initially, the PDGF family consisted only of PDGF-A and -B, but in 2000 and 2001, two more ligands were unexpectedly discovered. These novel ligands were named PDGF-C and PDGF-D, and added new complexity to PDGF signaling, which is still not completely understood. Moreover, the biological function of the latest member in the PDGF family, PDGF-D, is obscure. The aim of the work described in this thesis was to investigate the biological role(s) and functional relevance of PDGF-D, both during physiological and pathological conditions, and thereby enhance the understanding of PDGF signaling.

This introductory chapter will mainly focus on what is known about the structure, regulation and function of the PDGFs and their receptors in physiological and pathological settings. It will also introduce NRP1 and its known interactions with PDGFs, as this thesis describes a novel role of NRP1 as a co-receptor for PDGF-D.

1.1 THE PDGF SYSTEM

The PDGF family of growth factors consists of four ligands, PDGF-A, -B, -C, and -D, and two receptors PDGFR α and PDGFR β (2). The PDGFs were discovered already in the 1970's, and were among the first growth factors to be characterized, and the PDGF/PDGFR signaling have served as a model system for other growth factor families signaling through tyrosine kinase receptors, and is one of the most well-studied ligand-receptor systems.

1.1.1 PDGF ligands, structure and activation

The four PDGF ligands belong to the vascular endothelial growth factor (VEGF)/PDGF superfamily, which all share a conserved growth factor domain (GFD) with eight conserved cysteine residues that are important for disulphide bond formation and protein structure (3). Grouping by homology divides the PDGFs in two structural subgroups, also corresponding to the order in which the PDGFs were discovered. Thus, PDGF-A and PDGF-B form one subgroup where their N-terminal signal peptide is followed by a pro-peptide, the GFD and a C-terminal basic retention motif that allows for interactions with the extracellular matrix. PDGF-C and PDGF-D form a second subgroup as they have an additional N-terminal CUB domain (named after the first three proteins it was found in; Complement subcomponents C1r/C1s, Urchin EGF-like protein and Bone morphogenic protein 1), followed by a hinge region and the GFD, but no C-terminal retention motif (reviewed in (2)). The CUB-domains block receptor binding of the GFD, and need to be proteolytically removed before receptor binding and signaling can be induced. CUB domains are generally found in extracellular and membrane-associated proteins involved in developmentally regulated processes and are thought to mediate extracellular binding (reviewed in (4) and in (2)).

The PDGF ligands are encoded by four separate genes, which are located on different chromosomes. The *Pdgfd* gene is very large and spans over 200.000 base pairs, half of which are in the massive first intron. It is located on chromosome 11q22.3 in humans, and on 9A1 in the mouse (5-7) and is comprised of seven exons. Exon 1 encodes the signal peptide, exon 2-3 the CUB domain, exon 4 the hinge region between the two structural domains, and lastly, exon 6-7 encode the GFD (6, 8).

Biologically, the PDGFs occur as homo- or heterodimers (PDGF-AA, -AB, -BB, -CC, and -DD), which all require proteolytic processing before they can bind to their receptors. PDGF-A and -B are processed and activated prior to secretion, while PDGF-C and -D are activated extracellularly (reviewed in (2)). In PDGF-A and PDGF-B, the N-terminal propeptides are cleaved off intracellularly before secretion, by the dibasic-specific proprotein convertase furin (reviewed in (2)). In contrast to PDGF-A and PDGF-B, PDGF-C and PDGF-D are secreted in their full-length form, in which the CUB domains prevent receptor binding and needs to be cleaved off before receptor binding and signaling can occur. The extracellular cleavage is performed by serine proteases. PDGF-C is cleaved by tissue plasminogen activator (tPA), while PDGF-D is cleaved by urokinase plasminogen activator (uPA) and matriptase. However, there might be other proteases capable of cleaving PDGF-C and PDGF-D that are still unknown. The cleavage site for PDGF-D is located in the hinge region (R247/R249) between the CUB and GFD domains (9-11).

In addition to their full-length forms, PDGF-A and PDGF-B also exist in isoforms without the C-terminal retention motif; PDGF-A has an alternative splicing form, encoded by exons 1-5 and the short (3 amino acids) exon 7, while in PDGF-B, the retention motif can be removed by post-translational or extracellular processing (12-15). No additional isoforms have been reported for PDGF-C, but several splicing forms have been reported for PDGF-D. In human, the main isoform consists of 370 amino acids, and the full-length peptide chain had a calculated molecular weight of ~40.3 kDa migrates on SDS-PAGE as a M_r 90 kDa dimer, or as a M_r 55 kDa monomer (5). The cleaved, activated dimer migrated at M_r 35 kDa, and the cleaved, activated monomer at M_r 20 kDa (6). PDGF-D has a second splice variant of 364 amino acids, lacking 6 amino acids in the region upstream of the CUB-domain (5), and in mouse, a third splicing variant has been reported, which is lacking exon 6 and has no mitogenic properties in the GFD (16). Little is known about these alternate splicing forms of PDGF-D, and therefore, only the first isoform will be discussed in this work.

1.1.2 PDGF receptors, structure and signaling

PDGFs act by binding to the tyrosine kinase receptors PDGFR α and PDGFR β , which induces downstream signaling and thereby modulates cellular functions. The PDGF receptors form dimers upon ligand binding, with different specificity for the ligands. PDGF-AA, -BB, -AB and -CC all bind to the PDGFR α homodimer and PDGF-BB and -DD bind to the PDGFR β homodimer (reviewed in (2)). Each receptor monomer has five extracellular immunoglobulin (Ig)-like domains, whereof the three N-terminal ones are involved in ligand binding, followed by a trans-membrane domain and an intracellular split tyrosine kinase domain (reviewed in




Figure 1. PDGF processing and secretion. PDGF-A and PDGF-B propeptides are cleaved off intracellularly by furin and other proprotein convertases. PDGF-A_{long} and PDGF-B are kept close to the cell of origin through ECM binding of their C-terminal retention motifs. PDGF-A_{short} is diffusible. The PDGF-B C-terminal retention motif can be cleaved off to yield a diffusible PDGF-B. PDGF-C and PDGF-D are secreted in their inactive, diffusible forms, and require proteolytical removal by serine proteases of their CUB-domains to become activated.

(1)). Binding of a PDGF ligand to a receptor monomer induces receptor dimerization, whereupon conformational changes lead to auto-phosphorylation of the intracellular tyrosine residues and enables binding of SH2 (Src Homology 2) domain signaling molecules. Structural differences between the receptor monomers affect the binding site locations and affinities for SH2, and thus also the downstream signaling.

The PDGF receptor dimers formed upon ligand binding have different specificities for the ligand dimers, and while PDGF-AA, -BB, -AB and -CC all bind to the PDGFR α homodimer PDGF-BB and -DD bind to the PDGFR β homodimer (reviewed in (2)). All ligand dimers, except PDGF-AA, can bind to the PDGFR $\alpha\beta$ heterodimer *in vitro*, although a biological relevance of this receptor dimer has not yet been established in physiological conditions (reviewed in (2)). However, PDGFR $\alpha\beta$ dimers are present in certain tumors, and it is possible that also PDGF-DD signals through the heterodimeric receptor complex under pathological conditions (6, 17).

Activated PDGFRs interact with different families of SH2-domain containing molecules; some have enzymatic activity (Src, RasGAP, SHP-2, PLC- γ), others are adaptor molecules (Grb2, Nck, Shc, Crk, Alix and subdomains of phosphatidylinositol 3'-kinase (PI3K)) leading




Figure 2. PDGF receptor specificity. All PDGFs except PDGF-D bind to PDGFR α , but only PDGF-B and PDGF-D bind to PDGFR β . PDGF-AB, PDGF-BB, PDGF-CC and PDGF-DD can bind to the PDGFR $\alpha\beta$ heterodimeric receptor, but it is still unclear whether this receptor has a physiological relevance.

to downstream signaling. PDGFRs also bind signaling enhancers (NHERF), and activate transcription factors (STAT family) (reviewed in (1) and in (18)). An important PDGFR-induced pathway is the diverse PI3K pathway, which mediates actin re-organization and migration, as well as proliferative and anti-apoptotic responses, through Akt-mediated activation of the transcription factor NF- κ B. Other responses induced by PDGFRs are PLC- γ -mediated Ca^{2+} mobilization, and Ras and Src mediated activation of the Myc and Erk/MAP kinase pathways that promote proliferation (reviewed in (1)). The signaling outcome also depends on target cell-specific intracellular signaling pathways, and crosstalk between different pathways, which in some cases even counteract each other (reviewed in (1)). Most pathways are induced by all three PDGFR dimers, but the PDGFR β intracellular domain is more potent in inducing migration, and also has unique signaling capacities needed for mural cell function (19).

Following PDGFR activation and adaptor protein binding, receptors are removed from the cell surface by endocytosis. Vesicle forming proteins named clathrins are recruited to the PDGFR cytosolic domains to make them accumulate in clusters, which are subsequently reabsorbed into the cell and sorted for recycling or lysosomal degradation (reviewed in (20) and in (18)).

1.2 FUNCTIONAL ASPECTS OF PDGF SIGNALING

The PDGF system have been extensively studied *in vitro* and *in vivo*, and is known to be of vital importance during development, as both PDGFR null mice phenotypes are embryonic lethal. PDGFR α signaling is indispensable in organogenesis, and PDGFR β signaling is crucial for the development of the vascular system. The PDGFRs are expressed in cells of mesenchymal origin, such as fibroblasts, smooth muscle cells and pericytes (a perivascular supportive cell type), which are often found in connective tissues, while ligands are mostly expressed in neighbouring cells. Knowledge of the sites of expression for each of the effectors is necessary when trying to understand a functional role, as this often reflects sites of biological implications. Other important parts of the regulation include proteolytic activation of the ligands, bioavailability and interactions of the ligands with the extracellular matrix (ECM).

1.2.1 Expression and regulation

PDGFR α , PDGF-A and PDGF-C are expressed already in early embryogenesis, and in greater abundance than the more vascular PDGFR β , PDGF-B and PDGF-D during the early stages of development (reviewed in (2)). Both PDGFRs are expressed by mesenchymal cells, PDGFR α particularly by certain subtypes of progenitor cells (lung, skin, intestine and oligodendrocyte progenitors) while PDGFR β is strongly expressed in vascular smooth muscle cells (vSMCs) and pericytes throughout vascular system (reviewed in (2)). PDGFR β is also often used as a pericyte marker in adult tissue (23). It is noteworthy that no physiological function for the PDGFR $\alpha\beta$ heterodimer has been identified, and that PDGF and PDGFR expression levels are lower in adult (reviewed in (24)). The detailed expression of the ligands is complex, but in general, PDGF-B is expressed by endothelial cells, megakaryocytes and neurons while PDGF-A and PDGF-C are expressed in epithelial cells, muscle, and neuronal progenitors with a partial overlapping expression pattern. Their expression patterns have been reviewed in detail (2).

The expression pattern of PDGF-D, the most recently found PDGF family member, is not as well characterized as for the other PDGFs. However, *Pdgfd* messenger ribonucleic acid (mRNA) is known to be present in most adult tissues in both human, rat and mouse (5, 6, 25) further discussed in Paper I). The cellular sources of PDGF-D are not very well described, probably as a consequence of lack of both good commercial antibodies and *in situ* hybridization probes against human and especially mouse PDGF-D. Moreover, until our recent publication of the *Pdgfd*^{-/-} mouse strain with a *LacZ* reporter gene expressed under the *Pdgfd* promoter, reporter mice have also been lacking (see Paper I (26)). In spite of the lack of good commercial tools for histological detection of PDGF-D, there are a number of reports of PDGF-D expression. These observations are summarized below, in relation the expression of PDGFR β and PDGF-B.

Similar to PDGF-B and PDGFR β , PDGF-D is expressed at low levels in the normal adult vasculature, although there have been some discrepancies regarding its exact localization.

PDGF-D expression has been reported in all three vascular layers (endothelium, tunica media, tunica adventitia), by endothelial cells, vSMCs and fibroblasts (8, 27, 28). During development, PDGF-D and PDGFR β are co-expressed in arterial vSMCs, and both are present in blood vessels lining the vertebra, as well as in skin, skeletal muscle, liver, and lung (29). The same study also reported PDGF-D mid-gestational expression in epicardial and endocardial cardiomyocytes, and in myocardium, whereas cardiac PDGFR β was mainly observed in the vasculature (29). In the developing avian heart, PDGFR β expression has been reported also in the atrio-ventricular cushions that will develop into cardiac valves and connect to the septa, while PDGF-B was seen in endothelial cells, vSMCs, nerves and in the developing ventricular septum (30).

In the adult heart, PDGF-B and PDGFR β levels are very low, but PDGFR β is present in vSMCs (reviewed in (24)). Presence of PDGF-D has been reported in adult myocardium (29). In the kidney, PDGFR β is expressed by glomerular mesangial cells and vSMCs, while PDGF-B is expressed by endothelial cells. In developing and adult kidney, PDGF-D expression is present in multiple cell types of mesenchymal origin, such as arterial vSMCs and fibroblasts, both in mouse and human. PDGF-D is also expressed in neighbouring cells, such as glomerular podocytes and other epithelial cells of the nephron (5, 31, 32). The glomerulus was recently reported to display a species-specific expression pattern; in human, PDGF-D expression was seen in podocytes but not in mesangial cells (specialized glomerular pericytes), whereas the opposite was seen in the mouse (32). In the anterior part of the eye surrounding the lens, PDGF-D expression has been reported in the epithelial layers of the ciliary body and iris, both during development and in adult tissue. PDGFR β was expressed in the epithelium of the lens and cornea and in the mesenchyme surrounding the optic cup during development (33, 34). There are also reports on hematopoietic stem cells expressing PDGF-D, and adipose-derived stem cells co-expressing PDGF-D and PDGFR β but not PDGF-B (35, 36). PDGF-D expression has also been observed in the rat central nervous system, both during development (E16) and in adult tissue (37).

In summary, PDGF-D is present in both developing and adult tissues, where it is expressed both in cells of mesenchymal origin (vSMCs, mesangial cells and fibroblasts) and in neighboring cells, such as endothelium and certain epithelial tissues. This expression pattern suggests that PDGF-D can act through both paracrine and autocrine modes of signaling. Recently, we published a report on a *Pdgfd*^{-/-} mouse strain, carrying a *LacZ* reporter gene inserted into the PDGF-D gene to characterize the global and vascular expression of PDGF-D in detail (26). Our findings are further discussed in Paper I.

Regulation and modulation of PDGF signaling

As discussed above, the downstream signaling pathways appear to be similar between the two PDGFRs, while the expression of both receptors and ligands appear to be dependent on spatial and temporal regulation together with tissue specific environmental factors.

During physiological conditions, PDGFs mainly act by paracrine signaling, and consequently, the bioavailability of the ligands and receptors in different tissues is a major regulatory aspect, which is in turn depending on a multitude of regulatory mechanisms. A major regulatory factor for PDGF-C and PDGF-D is proteolytic activation that is needed to enable receptor binding (discussed in section 1.1.1). The proteolysis is regulated through local expression and activation by tPA, uPA and matriptase, and possibly also by other unknown proteases (9, 11, 38). Interactions with extracellular elements serve to prevent diffusion and/or receptor activation, and also mediate establishment of signaling gradients that are important for chemotaxis (reviewed in (39)). The retention motif bearing isoforms of PDGF-A and PDGF-B bind to heparan sulfate, and in blood, PDGF-B availability is also regulated through binding of the plasma protein α_2 -macroglobulin. It is not clear to what extent the latently secreted PDGF-C and PDGF-D interact with ECM through their CUB-domains (reviewed in (2)). However, unpublished observations from our lab indicate that full-length PDGF-D is more diffusible than the cleaved, activated growth factor. Signaling modulation through crosstalk with other signaling pathways has been described between PDGFR β and the adhesion receptor integrin $\alpha_v\beta_3$ (reviewed in (1)), thus indicating interference with or from cell-cell or cell-ECM contact. Modulation by co-receptors can act to block, enhance or alter intracellular signaling pathways or affect receptor availability. For PDGFs, this type of interactions have been reported together with low-density lipoprotein receptor related protein (LRP) (40), the uPA receptor (uPAR) (41) and Neuropilin 1 (NRP1) (42, 43). The role of NRP1 in modulation of PDGFR β signaling will be further discussed below and in Paper III. Receptor availability can be modulated through receptor trafficking upon endocytosis. For PDGFR β , endocytosis has been reported to be modulated by LRP (44) and Eph/ephrin signaling (45). Moreover, the metalloproteinase ADAM10 has been identified as a sheddase that cleaves off the extracellular domain for PDGFR β , although a functional role of PDGFR β shedding has not yet been found (46). Receptor autoactivation upon increased levels of reactive oxygen species (ROS) can induce PDGFR β -mediated proliferation and migration (47, 48).

1.2.2 Physiological roles

A substantial part of the knowledge of the basic PDGF functions are derived from *in vitro* studies, in which the PDGFs have been shown to mediate survival, proliferation and migration. There have also been extensive studies of gene expression, using transgenic mouse models and different loss- or gain-of-function strategies providing a deeper understanding of many complex physiological functions *in vivo*. During embryonic stages, PDGFs act to guide proliferation, recruitment and migration of cells, while in adult tissues, they mainly act to maintain homeostasis. In this section, the roles of PDGFs in development and adult tissues are discussed, with focus on the vascular function, as this is the main site of expression for PDGFR β , the signaling receptor of PDGF-D

PDGFs during development

Although *in vitro* studies have shown that the downstream signaling pathways between PDGFR α and PDGFR β largely overlaps, ablation of the two receptors give rise to very different phenotypes (reviewed in (49)).

PDGFR α and PDGF-A ablation gives rise to similar, multifaceted, phenotypes, with defects in oligodendrocytes, chondrocytes, neural crest, alveolar SMCs, kidney fibroblasts, chorioallantoic plate, Leydig cells, intestinal mesenchyme and dermis. *Pdgfa*^{-/-} animals die perinatally due to respiratory problems, while *Pdgfra*^{-/-} embryos die already at mid gestation (reviewed in (49)). The PDGF-C phenotype is background dependent; *Pdgfc*^{-/-} mice on a 129S1/Sv genetic background die perinatally from feeding and respiratory problems due to a cleft palate, whereas *Pdgfc*^{-/-} mice on a C57BL/6 genetic background survive, but display cerebral ventricular malformations, abnormal vascularization and skeletal deformations (50, 51). Loss of both PDGF-A and PDGF-C phenocopies the developmental defects in *Pdgfra*^{-/-} mice (50).

The *Pdgfrb*^{-/-} and *Pdgfb*^{-/-} phenotypes both gives rise to a general loss of vSMCs, and embryos die due to internal bleeding around birth with a slightly worse phenotype in *Pdgfrb*^{-/-} animals. Both *Pdgfrb*^{-/-} and *Pdgfb*^{-/-} animals also display abnormal cardiac innervation (52, 53). Moreover, both PDGFR α and PDGFR β are needed for cardiac neural crest development (reviewed in (2)).

Given the similar phenotypes of the *Pdgfrb*^{-/-} and *Pdgfb*^{-/-} mice, PDGF-D knockout mice were not expected to have a severe phenotype, at least not during embryonic development. However, there are some slight deviations between the *Pdgfrb*^{-/-} and *Pdgfb*^{-/-} phenotypes; PDGFR β -positive pericyte progenitors are present in skeletal muscle, skin and the adrenal gland in the less severe *Pdgfb*^{-/-} mouse phenotype (reviewed in (1)). It has been suggested that compensational PDGF-D signaling could be responsible for these phenotypic differences.

Little is known about the developmental role of PDGF-D; expression data indicate the presence of PDGF-D in cardiac, renal and cerebellar tissues, and a role in lens formation has been suggested (29, 31, 33, 37). Nevertheless, ligands need receptors to exert their functions, and with PDGFR β as the only known receptor for PDGF-D, their functional roles are expected to be coherent. PDGFR β is mainly expressed in vSMCs/pericytes and thus, PDGF-D is also expected to have a functional role for these cells.

The role of PDGFR β in angiogenesis and vascular maturation has been extensively studied using different mouse models and reporter genes, leading to the understanding that PDGF-B signaling through PDGFR β is crucial for proper pericyte recruitment and attachment to the vasculature (54). During angiogenesis, the endothelium adopts a more active phenotype to build new vessels. PDGF-B is secreted from endothelial *tip cells* at the angiogenic front to attract PDGFR β -expressing pericytes or mural cell progenitors that migrate along the newly formed vessel sprouts (54). The *tip cells* are VEGF-sensitive and instruct neighboring

endothelial cells to take on the proliferative *stalk cell* phenotype through Notch signaling, and thereby promote to elongation of the vascular sprouts (55). In the newly formed endothelium, PDGF-B signaling through PDGFR β promotes maturation and stabilization. Moreover, their signaling also controls mural cell fate and proliferation (56, 57). Consistently, PDGF-B expression is strong also in growing arteries, where mural cells are actively recruited (23, 54). Later on, and in the adult stage most endothelial cells display a third, resting *phalanx cell* phenotype (58). Here, stabilizing pericyte-to-endothelial cell signals regulate the endothelial proliferation, survival, migration and differentiation, thereby repressing unwanted angiogenesis. Vessels that lack pericytes are leaky and have increased capillary diameter. The endothelial cells in such vessels display hyperplasia and excessive folding of the luminal membrane, accompanied by altered expression of junction proteins (59). Thus, pericytes act as negative regulators of endothelial proliferation, and also instructs the endothelium to keep a smooth luminal surface membrane (reviewed in (2)).

As described in section 1.2.1, many stem and progenitor cells also express PDGFs during development, and their activity is often regulated by PDGF signaling from neighbouring cells. One example is in the Sertoli cells of the testis, where both PDGF-B and PDGF-D are expressed and regulate proliferation and migration of spermatogonial cell precursors (60).

Physiological role in adult

The embryonic lethality of *Pdgfra*^{-/-} and *Pdgfrb*^{-/-} animals has lead to a focus on the developmental functions of these receptors. The PDGFRs are expressed in lower levels also in adult, dormant tissues (reviewed in (24)), and it has been suggested that the role of PDGF signaling is less important in the adult physiology than in the developmental. Consistently, pharmacological blocking of PDGFR signaling by the tyrosine kinase inhibitor Imatinib (also targeting c-kit and Abl) is tolerated relatively well (61). Imatinib treatment did, however, give adverse effects from the gastrointestinal system, accompanied by myalgia and edema, indicating sites of PDGFR activation in the adult (61). Indeed, PDGFR α is expressed in the gut, and PDGFR β -expressing perivascular cells control vessel integrity and interstitial fluid pressure (62). Interestingly, a PDGF-D neutralizing monoclonal antibody (CR002) has been evaluated in a Phase I trial. Antibody-mediated amelioration of proteins produces few side effects, and is as close to a knockout that one can get in the human situation (63).

Nevertheless, PDGFs are needed in adult tissue homeostasis, and mouse models of partial, transgenic and conditional loss or gain of function have contributed to the understanding of PDGF signaling in adult settings. One important model is the *Pdgfb*^{ret/ret} mouse strain that lacks the PDGF-B retention motif, which normally retains the secreted ligand close to the endothelial cell, as a chemotactic signal for vSMCs. The *Pdgfb*^{ret/ret} mice survives until adult age, but are smaller than its wildtype littermates, and suffers from a vascular phenotype leading to renal and retinal dysfunctions. This model has also shown that the PDGF-B retention motif is necessary for proper pericyte recruitment and attachment to the capillaries (64). Further studies of the *Pdgfb*^{ret/ret} mouse strain has also lead to the understanding that

pericytes are needed to maintain the blood brain barrier, and that this is regulated through PDGF signaling (65).

As implied by the name, PDGFs were originally isolated from platelets, where they play an important role in wound healing. Platelets store PDGF-A, -B, and -C and other growth and clotting factors in α -granules, which are released upon tissue damage, in order to prevent blood loss and maintain homeostasis (66, 67). PDGF signaling is needed throughout the wound healing process, and contributes to a number of different events including inflammation, formation of granulation tissue, re-epithelialization, remodeling and neovascularization ((66) and reviewed in (1)). PDGF-B has been shown to be especially potent and is released from platelets immediately upon wounding, and stimulates the healing process through increased fibroblast proliferation, ECM production and neovessel formation (reviewed in (2)). Recombinant human PDGF-B (Becaplermin) is used in the clinic to enhance wound healing (68). In similarity to PDGF-B, PDGF-D also serves in recruitment and proliferation of monocytes/macrophages and vSMCs (69) and stimulates secretion of ECM, as well as matrix degradation factors (MMPs/TIMPs) by vSMCs and fibroblasts (70). Part of the wound healing process is also neovessel formation, where PDGFs induce secretion of VEGFA to promote angiogenesis, which is also guided through hypoxia and ECM scaffolding (reviewed in (71)). Other cases when adult physiological angiogenesis is needed include the adaptive response of cardiac and skeletal muscle upon physical exercise, and in the uterine cycle, where the tissue is shed and rebuilt in a cyclic manner.

Another function of PDGFR β -expressing mural cells is in control of interstitial tissue pressure, through maintaining vascular impermeability and contraction of ECM, which modulated by both PDGF-B and PDGF-D signaling (69, 72).

PDGFs regulate mesenchymal stem cell functions also in adult tissues. During the uterine cycle, PDGFR β -expressing cells found in perivascular locations have been suggested to be responsible not only for angiogenesis but also for the cyclical growth of the endometrium (73). A special function in certain stem cells is also seen for PDGF-D. In contrast to PDGF-B, PDGF-D is occasionally co-expressed with PDGFR β , thus facilitating self-maintaining autocrine signaling loops that promote proliferation and migration, as seen in adipose tissue derived stem cells (74).

In summary, as a PDGFR β ligand, PDGF-D may induce responses that overlap with those of PDGF-B, such as survival, proliferation, migration, vascular maturation, and regulation of interstitial fluid pressure and ECM homeostasis. Thus, there may be some functional redundancy in PDGFR β signaling between PDGF-D and PDGF-B. In consistence, we recently reported our finding that the *Pdgfd*^{-/-} mice display no major phenotype and survive to adulthood. However we also reported that PDGF-D appear to be involved in mural cell maintenance during physiological conditions, which is further discussed in Paper I (26).

1.3 PATHOLOGICAL ASPECTS OF PDGF SIGNALING

Dysregulation of PDGF activity is seen in a number of common pathological conditions, such as fibrosis, atherosclerosis and cancer. To better understand the role of PDGF signaling in these conditions, extensive studies have been performed, including different overexpression strategies *in vivo*. These studies indicate that PDGFs have strong impact on the progression of these pathologies, through promotion of excessive proliferation, ECM synthesis and migratory signals, commonly affecting different types of SMCs and/or fibroblasts.

1.3.1 Fibrosis

Fibrosis is a common condition that is mechanistically similar to exaggerated wound healing, but also occurs in response other types of tissue damage. The pathological process is characterized by excessive deposition of collagens and other ECM components in the interstitium, leading to scarring and disruption of tissue function. These events are also firmly connected to transforming growth factor β (TGF- β) signaling, which is a very potent inducer of ECM secretion (75). Thus, a central player in fibrosis is the fibroblast, a heterogeneous cell type that varies between organs, which responds to TGF- β , PDGF, and other stimuli (reviewed in (76) and in (2)).

PDGFs contribute to the fibrotic process by proliferative and chemotactic signaling, especially through myofibroblast expansion (reviewed in (77) and in (2)). All four PDGFs are upregulated during fibrosis, and cause massive fibrotic responses when overexpressed in animal models (reviewed in (2)). High PDGF expression has been described in pulmonary, hepatic, renal and cardiac fibrosis, although the expression and regulation of each PDGF ligand appears to be organ-specific also in fibrosis (reviewed in (77)). Interestingly, crosstalk has been reported between PDGFs and the pro-fibrotic TGF- β , which seems to contribute to the spatial regulation of the different ligands. In the case of PDGF-D, TGF- β has been shown to promote signaling in cardiac fibroblasts, but inhibits PDGF-D expression in the lung (78, 79). The pro-fibrotic potential of PDGF-D in heart and kidney has been demonstrated by overexpression studies *in vivo* (29, 32, 80), and PDGF-D/PDGFR β signaling is also implicated in liver fibrosis (81-83). PDGFR β -mediated fibrotic responses are executed by fibroblasts, myofibroblasts and specialized pericytes, such as hepatic stellate cells and renal mesangial cells that have been activated into a fibroblast-like state. Like other pericytes, they express PDGFR β (84), and PDGF-D has been shown to exhibit mitogenic and fibrogenic effects on these cells (81, 83, 85, 86). Similar to the other PDGFs, PDGF-D contributes to accelerated ECM deposition. In both hepatic stellate cells and cardiac myofibroblasts, PDGF-D has been shown to upregulate tissue inhibitor of metalloproteinase (TIMP)-1, which attenuates matrix metalloproteinase (MMP) activity, thereby decreasing ECM degradation (78, 85). Notably, PDGF-D-mediated ECM remodeling is implicated in epithelial-to-mesenchymal transition (EMT) in cancer (87), a process that is activated also in fibrosis and chronic inflammation, (reviewed in (88)). Thus, it is not unlikely that PDGF-D contributes to EMT also in fibrosis, but further studies are needed on this subject.

In severe fibrosis, the irreversible tissue scarring leads to loss of tissue function, and therefore early anti-fibrotic/anti-inflammatory treatments are used. Several studies suggest PDGF-D as a suitable therapeutic target in kidney fibrosis, and an anti-PDGF-D monoclonal antibody (CR002) has been shown to prevent renal fibrosis in mice (82, 86, 89). Also, the *Pdgfd*^{-/-} mice (presented in Paper I) show reduced renal fibrosis upon experimental induction of renal scarring (32).

1.3.2 Atherosclerosis and other vascular pathologies

Cardiovascular disease is a major cause of death in the western world. A major risk factor is atherosclerosis, a process in which lipid-containing plaques are slowly formed in the vascular wall. With time, the plaques may rupture and cause thrombosis or internal bleedings that can be life-threatening. Atherosclerotic plaques appear in vessel areas exposed to low shear stress, turbulent flow and oscillating flow (reviewed in (24)). Plaques are formed over decades, initially through lipid accumulation, followed by endothelial dysfunction and infiltration by macrophages and other immune cells into the vascular wall, leading to chronic inflammation.

In contrast to the healthy vasculature that expresses only low, or undetectable levels, of PDGFs, all four PDGF ligands are present in the atherosclerotic vessel wall, and are mainly secreted by endothelial cells, vSMCs and macrophages. Also, both PDGF receptors are upregulated, and expressed by vSMCs and macrophages ((28, 70) and reviewed in (24)). PDGFR β expression is stronger than that of PDGFR α , and PDGFR β signaling has been shown to be a driving force in the atherogenic process through chemokine signaling that induces leukocyte migration (90). In the later stages, active vSMCs play a central role, as they proliferate and contribute to the thickening of the vascular wall (reviewed in (91)). Atherogenic stimuli (ECM, cytokines, shear stress, reactive oxygen species and lipids) promote the switch from a contractile to synthetic phenotype of vSMCs, which is also associated with increased PDGFR β expression and signaling (reviewed in (92)). PDGF-B and PDGF-D are both upregulated in endothelial cells exposed to atheroprone blood flow, and promote the vSMC phenotypic switch ((93) and reviewed in (24)). For PDGF-D, this is mediated through upregulation of the differentiation repressor gene Kruppel-like factor-4, and downregulation of contractile proteins, such as α -smooth muscle actin (α SMA) and the smooth muscle myosin heavy chain (93). PDGFR β , PDGF-B and PDGF-D also are induced during monocyte-to-macrophage differentiation, and they are all strongly expressed in fatty streaks, where PDGFR β signaling has a strong impact on monocyte/macrophage migration (28, 70). Moreover, PDGFs promote stabilization through ECM synthesis, as well as ECM remodeling through MMP and TIMP expression (27, 70).

Notably, recent studies have reported genetic variations in the form of single-nucleotide polymorphisms (SNPs) in the *PDGFD* gene to be associated with coronary artery disease (94), and non-hypertensive intra-cerebral hemorrhage (95). This indicates that aberrant PDGF-D signaling could be causative in atherogenesis and cardiovascular disease, emphasizing the importance of PDGF-D in these conditions.

PDGFs in response to vascular injury, rejection and restenosis

Vascular injury, restenosis and chronic rejection gives rise to conditions that resembles atherosclerosis. Upon vessel injury, medial vSMCs migrate from to the intima, where they proliferate and switch to a synthetic phenotype in a PDGFR β dependent manner, resulting in neointimal hyperplasia. PDGF-D is upregulated in the intima following angioplasty, and is also thought to contribute in this process, next to PDGF-B ((27) and reviewed in (24)). The response to chronic cardiac allograft rejection also manifests as a vascular disease, and is histologically characterized by concentric luminal stenosis (vascular narrowing) (reviewed in (96)). In rat cardiac allografts, PDGF-D, but not PDGF-B, was induced, and increased the pro-fibrotic and pro-arteriosclerotic responses leading to chronic rejection, through the TGF- β 1 pathway (97).

PDGF signaling is of importance also in vascular pathologies involving the blood brain barrier integrity in the central nervous system. In stroke, PDGF-C signaling through PDGFR α is involved in the acute opening the blood-brain barrier (98, 99). In the chronic disease named amyotrophic lateral sclerosis (ALS), high neuronal expression of PDGF-C leads to disruption of the blood-spinal cord barrier, and contributes to the early onset of the disease (100).

1.3.3 Cancer

In cancer, malignant cells grow in an uncontrolled manner and disrupt normal tissue functions. The six hallmarks of cancer established by Hanahan and Weinberg include proliferative signaling, evading growth suppressors, resisting cell death, enabling replicative immortality, inducing angiogenesis, and activating invasion and metastasis (101). The tumor is built up of cancer cells, some of which have stem cell properties, and others have invasive capabilities. The tumor microenvironment also contains stromal cells, which include cancer-associated fibroblasts (CAFs), immune inflammatory cells, endothelial cells and pericytes as well as local and bone marrow-derived stromal stem and progenitor cells (reviewed in (102)).

The first connections between PDGFs and cancer arose in 1983, when PDGF-B was sequenced and proved to be similar to v-sis, the transforming protein of simian sarcoma virus (103, 104). Today, all PDGFs and PDGFRs have been connected to different tumor types, and are known to be capable drivers of tumorigenicity through several of the cancer hallmark functions; proliferation, angiogenesis and migration (reviewed in (18)).

PDGF ligand overexpression is seen in carcinomas, tumors of endodermal or ectodermal origin, where they promote tumorigenesis and metastasis through recruitment and stimulation of stromal mesenchymal cells in a paracrine manner. PDGFs can act through autocrine signaling loops in cases where the tumor cells express the receptor (reviewed in (105)). One example of autocrine signaling is in dermatofibroma protuberans, where PDGF-B is expressed under a collagen promoter, resulting in a COL1A1/PDGF-B fusion protein from which active PDGF-B is cleaved off (106).

Constitutionally active receptors are seen in tumors derived from PDGFR-expressing mesenchymal cell populations, such as sarcomas and glioblastomas, where they drive proliferation and clonal expansion through autocrine signaling (reviewed in (1)). These receptors are the result of gene translocations leading to fusion proteins as seen in hypereosinophilic syndrome, caused by a *FiP1L1-PDGFR α* fusion (107), and chronic myelomonocytic leukemia, caused by an *ETV6-PDGFR β* fusion (108).

Other types of genetic aberrations in PDGFs and PDGFRs have also been observed to cause cancer in the case of PDGF signaling. For PDGFR α , activating point mutations are commonly seen in gastrointestinal tumors (GIST) (109), as are gene duplications in glioblastomas (110). Overexpression of receptors can also lead to auto-activation (reviewed in (111)).

PDGFR β signaling in the tumor

In solid tumors, PDGFR β expression is mainly confined to stromal cells, such as CAFs, vSMCs and pericytes (reviewed in (112)), where PDGFR β is important in vascular stabilization. Tumor vessels are often leaky and fragile due to lack of supporting pericytes (reviewed in (102)). Both PDGF-B and PDGF-D, signaling through PDGFR β , stabilizes the vasculature through recruitment of pericytes (113), thereby enhancing vascular function and contributing to tumor growth. Pericyte PDGFR β signaling also provides survival signals to the endothelium, for example by upregulating the anti-apoptotic gene *Bcl2l2* (114). PDGFR β expression in CAFs is associated with aggressiveness and metastasis, and PDGFR β signaling in both CAFs and pericytes promote high interstitial fluid pressure in tumors through ECM interactions (115).

PDGF-D signaling through PDGFR β in cancer

PDGF-D has been reported to be a driver in tumorigenesis in different cancers, such as prostate and renal cell carcinoma, pancreatic adenocarcinoma, glioblastoma, schwannoma (derived from peripheral glia cells) and melanomas (reviewed in (116)). A major difference between the PDGFR β ligands, PDGF-B and PDGF-D, is that PDGF-D requires proteolytic activation, and thus, the enzymes that activate PDGF-D, uPA and matriptase, needs to be present in the microenvironment, as has been shown in prostate cancer (9, 11). The role of uPA in tumor growth, angiogenesis and metastasis is well studied (reviewed in (117)). For matriptase, a feedback loop has been reported, where the activated PDGF-D upregulates matriptase (118). Once activated, PDGF-D promotes proliferation, and stimulation of tumor cell growth (119). Similar to physiological conditions, PDGF-D and PDGF-B interact with the stroma through recruitment of PDGFR β -expressing cells (vSMCs and macrophages), and expression of VEGF and MMPs (120-124).

Coherently, gene expression profiling has shown that PDGF-D associates with genes active in cell adhesion, wounding, and immune system processes (17). Moreover, in dermis, non-malignant PDGF-D overexpression has been shown to increase interstitial fluid pressure (69). Both vascular stability (or impermeability) and high interstitial pressure act to impair drug

delivery, as the drug is maintained in the blood stream. Targeting of PDGFR β signaling improves drug delivery and efficacy of chemotherapy (125, 126). Inhibition of PDGF-D has been shown to attenuate growth, invasion and angiogenesis in a xenograft model of gastric cancer (127). However, in breast cancer PDGF-D blockade also promoted lymphatic metastasis by activation of CXCR4 (128).

PDGF-D in EMT, invasion and metastasis

PDGF-D and PDGFR β has also been implicated in invasion and metastasis, through the process of EMT, in which epithelial cells lose their apical-basal polarization and cell-cell adhesion to acquire a migratory, fibroblast-like phenotype ((129, 130) reviewed in (131)). During physiological conditions, embryonic EMT is essential for many developmental processes, and in adult tissue it is implicated in wound healing (reviewed in (131)) and in the pathological process of fibrosis (reviewed in (88)). In cancer, tumor cells reactivate the processes of EMT to become invasive and metastatic which is associated with upregulation of mesenchymal markers involved in invasion and migration (reviewed in (132)). PDGF-D appears to regulate the process of EMT on several levels, partially through the NF- κ B and Notch pathways (87, 121). This leads to downregulation of the epithelial adhesion protein E-cadherin, which in prostate cancer takes place through inhibition of microRNA-mediated regulation of the transcription factors Snail, Twist and ZEB1 (133-136). PDGF-D has also been shown to upregulate expression of MMPs and the cytoskeletal protein vimentin and other mesenchymal markers (124, 137), to promote degradation of basement membranes and ECM and thus also angiogenesis, metastasis and invasion.

PDGF-D and cancer stemness

There have been different theories on how tumors grow. Today, one that is favored by many researchers is the cancer stem cell (CSC) hypothesis. CSCs constitute a small subset of cells with the ability form new tumors when implanted into SCID mice, where they both self-renew and differentiate into more mature phenotypes. These cells only make up a small part of the total bulk of cancer cells, but are the main target to treat malignant and metastasizing disease (reviewed in (116) and in (102)). The origin of the CSCs is not clear, but they may be derived from adult stem cells, or more differentiated progenitor cells, that are present in normal tissues, and the source may also differ between tumors (reviewed in (102)). The traits of CSC and EMT-transdifferentiated cells overlap to a large extent, and therefore, PDGF-D signaling through PDGFR β has been suggested to be involved also in cancer stemness. It has been shown that PDGF-D transformed NIH-3T3 fibroblasts show CSC features, such as anchorage-independent growth in soft agar and ability to induce tumors in nude mice (119). Moreover, several studies have reported that high expression of PDGF-D induces EMT, and CSC-like capabilities in cancer cells, thus contributing to tumor aggressiveness (136, 138).

Recently, we published a study showing that in a RIP1-TAg2 model of neuroendocrine pancreatic cancer, the tumours in *Pdgfd*^{-/-} mice developed more slowly than those in *Pdgfd*^{+/+}

littermates. The possibility that this delay is because of altered CSC capacity is further discussed in Paper II.

1.4 NEUROPILIN 1

Neuropilin 1 (NRP1) and NRP2 are multifunctional trans-membrane co-receptors for members of the class-3 semaphorins (SEMA3) in neuronal axon guidance and vascular endothelial growth factors (VEGFs) in angiogenesis. However, interactions of NRP1 with ligands from other signaling systems, such as TGF- β 1, hepatocyte growth factor (HGF) and fibroblast growth factor (FGF) have been reported (reviewed in (139)).

There is a growing body of evidence showing that PDGFR signaling can be modified by NRP1 (42, 43, 140-144), and in Paper III, we provide evidence that NRP1 can act as a co-receptor for PDGF-D. In this section, NRP1 and its functions as a co-receptor for the structurally related members of the PDGF/VEGF superfamily are described.

1.4.1 Structure, function and expression of NRP1

The NRPs consists of seven sub-domains where the first five are extracellular; two CUB domains (a1 and a2), two coagulation factor V/VIII domains (b1 and b2) and a meprin, A5 μ -phosphatase domain (c). Following the trans-membrane domain, NRPs contain a short cytosolic tail with a PDZ (Post synaptic density protein, *Drosophila* disc large tumor suppressor, and *Zonula occludens-1* protein) binding domain (reviewed in (145)). Full-length NRP1 protein is present as a 130 kDa species, but there are also several splice variants that are less studied, and will not be discussed in this thesis (reviewed in (146)).

Ligand binding occurs at different sites of NRP1; SEMA3A binding requires the first three NRP1 sub-domains; a1, a2, and b1, whereas VEGF-A binds primarily to a negatively charged cleft in the b1 domain but also to the b2 domain (147, 148). VEGF-A₁₆₅ binds to NRP1 through a [KPRR] motif in the C-terminus, where the last arginine is crucial for binding (148, 149). The C-terminal arginine is shared by most of the NRP1-binding VEGF isoforms (reviewed in (146)). It has been suggested that NRP1 is present as homo- or heterodimers, but it is also possible that they oligomerize to form high-affinity binding sites for ligands (150).

NRP1 is expressed by a variety of cells, including vSMCs (143), endothelial cells, neurons, and different epithelial cells (reviewed in (151)). In the vasculature, NRP1 is mainly expressed by arterial endothelial cells, and during angiogenesis also by neural progenitors and macrophages. NRPs are also commonly expressed by tumor cells and tumor vasculature (reviewed in (152)). Studies of knockout and transgenic mouse models have shown that NRP1 is crucial for neuronal, cardiac and vascular development, and *Nrp1*^{-/-} animals die at E10-12.5, with VEGF-A/NRP1 signaling-dependent defects in angiogenesis, vascular branching and topographic origin of the coronary arteries (153-155). It is also clear that NRP1 expression requires tight regulation; when overexpressed, NRP1 resulted in a lethal (E12.5) phenotype with excessive capillary growth, hemorrhages, cardiac malformation and neuronal defects (156). Also, the cytoplasmic part of NRP1 is not needed for survival, as mice expressing a truncated NRP1 protein displayed a mild phenotype with increased retinal arteriovenous crossings, and reduced number of arteries in heart, kidney and skeletal muscle (157, 158).

In conclusion, the NRP1 co-receptor has a vital role in arteriogenesis and angiogenesis, and it is expressed in PDGFR β -expressing vSMCs and neighbouring endothelial cells. Thus, the correlating spatio-temporal expression of NRP1, PDGFR β and the PDGFR β ligands would facilitate their interaction.

1.4.2 NRP1-mediated modulation of signaling

Neuropilins are mainly modulators of signaling (reviewed in (146)), and appear to induce situation-specific responses. Enhancing functions are generally seen for SEMA3, VEGF-A, and PDGF ligands, while NRP1 has an inhibitory function for TGF- β . Interestingly, in angiogenesis, NRP1 enhances the endothelial tip cell phenotype through VEGFR2 signaling while it inhibits the stalk cell phenotype through inhibition of TGF- β R1/2, thus driving two different responses in the same cell type (reviewed in (159)). How NRPs exert their functions is still unclear, but suggested mechanisms include ligand capture, receptor inhibition, and competitive binding between ligands (160-162). The intracellular alteration of responses is mediated through enhanced receptor trafficking or enhanced cell migration or adhesion through direct or indirect interaction with integrins (141, 143, 163, 164).

Signaling occurs in both *cis* and *trans*. So far, signaling in *trans* has mainly been reported in tumor cells, where it counteracts angiogenesis by interference with VEGFR2 trafficking. NRP1 signaling in *trans* can also mediate adhesive functions, and thereby modulate the signaling time frame, or enhance cell-cell contact (165). In paper III, we show that PDGF-D induces complexes between NRP1 and PDGFR β , both in *cis* and in *trans*, which further extends the possibilities of modulation of PDGF-D signaling.

NRP1 in PDGF signaling

There is an increasing amount of data showing that NRP1 regulation of PDGF signaling appears to be important for vSMC function, and possibly also endothelial-mural cell communication (42, 43, 140-144). As discussed previously, the main PDGF ligand in the vasculature is PDGF-B, and several studies report that NRP1 can modulate PDGF-B-induced migration of vSMCs through tyrosine phosphorylation of the intracellular integrin adaptor protein p130Cas (141, 143, 166). NRP1 has also been reported to participate in PDGF-mediated recruitment and differentiation of MSCs into pericytes (140). Consistently, several studies have shown that impairment of NRP1 signaling disrupted PDGF-B-mediated vSMC migration (43, 142-144). NRP1 implications in vSMC-related diseases also appear to overlap with the indications for PDGFs; NRP1 contributes to neointimal hyperplasia, and promote the synthetic phenotype of vSMCs, through a pathway that is likely PDGF-dependent (144). NRP1 is also upregulated in response to PDGF-B, thus reinforcing NRP1-PDGF signaling (43).

Physical interactions between NRP1 and PDGFRs have been suggested, but the specific mechanisms for modulation of PDGF/PDGFR signaling by NRP1 are uncertain. Co-immunoprecipitation experiments have shown that NRP1 forms complexes with PDGFR α and PDGFR β in mesenchymal stem cells when stimulated with PDGF-A or PDGF-B, thus

indicating that NRP1 could act as a co-receptor for PDGFs (42). However, there have been contradictory reports on whether PDGF-B can bind to NRP1 (43, 143). These controversies may be explained by the presence of an additional, 250 kDa form of NRP1, which is glycosylated between the “b2” and “c” domains by chondroitin sulfate (CS) or heparan sulfate glycosaminoglycans. The CS-NRP1 enables cell-specific regulation of binding, and is present in vSMCs and certain tumor cells, where it has been shown to promote PDGF signaling. At the same time CS-NRP1 also inhibits VEGF-A binding to VEGFR2 (167).

We have found that PDGF-D is a much more specific ligand for the 130 kDa NRP1 species, than PDGF-B, and that PDGF-D binds NRP1 with about the same affinity as VEGF-A₁₆₅. This finding, and the role of NRP1 as a co-receptor for PDGF-D are discussed in paper III.

2 AIMS OF THIS THESIS

With the exception of PDGF-D, the PDGFs and PDGFRs have been well studied in different mouse models with genetic alterations. The aim of this thesis is to explore the physiological role of PDGF-D in health and disease, mainly by analyses of the *Pdgfd*^{-/-} mouse strain.

The specific aims include:

- To map the expression of *Pdgfd*, using the LacZ reporter gene in the *Pdgfd* knockout construct (Paper I)
- To investigate the phenotype the *Pdgfd*^{-/-} mice (Paper I)
- To evaluate the role of PDGF-D ablation in a genetic model of cancer (Paper II)
- To characterize the binding of NRP1 to PDGF-D, and the ability of PDGF-D to induce complex formation between NRP1 and PDGFRβ (Paper III)

3 PAPERS AND DISCUSSION

To highlight central findings and rationale, each paper will be summarized and discussed below. Methods and results are described in detail in the respective papers.

3.1 PAPER I – MICE LACKING PLATELET-DERIVED GROWTH FACTOR D DISPLAY A MILD VASCULAR PHENOTYPE

The roles of PDGF-A, -B and -C, and their signaling through PDGFRs have been studied through a large number of reporter gene, transgenic and knockout animal models, which have contributed greatly to the understanding of PDGF biology. However, the role of PDGF-D is still obscure, and prior to this study, there were no reports of reporter genes or knockout mice for PDGF-D.

In a ligand-receptor system such as the PDGF system, the phenotype of the ligands is expected to match that of the receptor, as seen for the *Pdgfra*^{-/-} and *Pdgfc*^{-/-} phenotypes compared to that of *Pdgfra*^{-/-}. The phenotypes of PDGF-B and PDGFRβ are very similar, and therefore, the phenotype of *Pdgfd*^{-/-} mice was expected to be milder. However, PDGF-D exerts its functions through PDGFRβ, which has a very distinct role in vSMCs/ pericytes, where it is needed for mural cell recruitment and proliferation during angiogenesis, and maintenance of the vascular homeostasis in the adult. PDGF-D expression has been reported in the vasculature previously (8, 27-29, 31), although there have been some inconsistencies in the reports of expressing cell types. PDGF-D has also been implicated in a number of vascular pathologies, and therefore, the vasculature was a candidate site of action.

In this study, we present the *Pdgfd*^{-/-} mouse strain, which is viable, fertile and show no gross abnormalities, in contrast to the other PDGF knockouts. The mouse has a normal life span, which enables future studies of PDGFRβ signaling in the adult settings. The study aim was to map the expression of *Pdgfd* through the *LacZ* reporter gene that was inserted in the targeted allele, and to explore the physiological functions of PDGF-D/ PDGFRβ signaling, with the vasculature as a starting point.

To identify organs of interest for PDGF-D function, we performed a general expression analysis on mRNA level, where adrenal gland, spinal cord, aorta, heart, uterus, cerebellum and lung were identified as high expression organs. These are all highly vascularized organs, and also, the vasculature was pointed out as a common expression site of *Pdgfd* between organs in our histology based reporter gene studies, thus supporting the notion that PDGF-D should have a vascular function. In concordance with previously published studies (27, 29), *Pdgfd* expression was seen predominantly in the endothelial compartment, but also occurred in vSMCs. We also confirmed that the *Pdgfd* expression was more prominent in arteries than in veins and frequently found in foci around vessel bifurcations. Moreover, there appeared to be a spatio-temporal regulation of expression, as it went from a strong, exclusively arterial expression at postnatal day (P4) to become more widely spread in adult vasculature, which could be representative of both endothelial-to-mural cell and mural cell autocrine signaling.

In total, our study confirms that PDGF-D can be expressed in both endothelial cells and vSMCs, and that the expression differs depending on vessel type and stage of life.

The other PDGFR β ligand, PDGF-B, is secreted by the endothelium in its active form but its expression is low or undetectable in the healthy, dormant endothelium. As PDGFR β is expressed in vSMCs, a possible function for PDGF-D could be in vascular maintenance or endothelial-to-mural cell signaling, where it would be activated upon need. Therefore, we examined vascular morphology and pericyte coverage. While we found no changes in endothelial appearance, nerve/glial antigen 2 (NG2)-expressing pericytes appeared disorganized and less attached in the cardiac vasculature of *Pdgfd*^{-/-} mice. We verified the downregulation of NG2 also on mRNA level by qPCR, and also found another mural marker, desmin, to be downregulated. To explore whether the deviations in NG2⁺ vasculature had implications on *Pdgfd*^{-/-} heart function, we performed a broad gene-expression analysis, including different immunological, developmental and vascular genes. We found a downregulation of *Gata4* and *Notch1* mRNA, two genes that regulate cardiac development and growth, cell fate, and maintenance of cardiac function in adult tissue, and thus, we continued the search of a cardiac phenotype. Clinical chemistry analysis revealed a modest increase in *Pdgfd*^{-/-} serum concentrations of Ca²⁺, and we observed a trend towards smaller hearts in the *Pdgfd*^{-/-} animals, compared to wildtype controls. Finally, we also detected a slight, but significant, elevation in both systolic and diastolic blood pressure compared to wildtype controls, suggesting a peripheral vascular or cardiac defect in *Pdgfd*^{-/-} mice.

As the *Pdgfd*^{-/-} animals survive, we did not expect to find any major heart defects. However, the slight increase in blood pressure indicates that something is different in the vasculature or the regulation of blood pressure, and it is tempting to speculate that it is related to the abnormalities seen in cardiac NG2⁺ pericyte morphology and/or serum Ca²⁺. It could also be connected to cardiac output, mean arterial pressure or peripheral resistance, which also affect the blood pressure blood pressure. It would be interesting to see whether anomalies are present also in other PDGFR β -expressing cells, such as epithelial cells, of the *Pdgfd*^{-/-} mice, and to assess cardiac output, and other cardiac parameters by echocardiography.

In summary, we present a *Pdgfd* knockout mouse strain with a mild vascular phenotype, thereby completing the PDGF “knockout family”. The *Pdgfd*^{-/-} mice will be a great aid in studying the physiological and pathological roles of PDGF-D/PDGFR β signaling, during both developmental and in adult stages of life. We provide new insights on both expression pattern and function of PDGF-D, suggesting a role in regulation of arterial blood pressure. We also describe a possible role for PDGF-D in mural cell physiology and regulation of vascular homeostasis.

3.2 PAPER II – FUNCTIONAL MALIGNANT CELL HETEROGENEITY IN PANCREATIC NEUROENDOCRINE TUMORS REVEALED BY TARGETING OF PDGF-DD

Tumors are heterogeneous, and consist of cancer cells of varying level of differentiation, and varying composition of the stromal compartment. PDGFs are commonly deregulated in

cancer, and do not only act on the tumor cells, but also on PDGFR-expressing stromal cells, such as CAFs, pericytes and vSMCs (reviewed in (112)), where they drive cellular processes such as proliferation, survival and motility, thereby promoting tumor growth and invasion (reviewed in (1)). Pharmacological blockade of PDGF signaling is routinely used for treatment of a diversity of malignancies (105). PDGF-D is upregulated in a number of different tumors, and is known to promote tumor progression through ECM remodeling and deposition. Lately, PDGF-D has also been reported to be involved in EMT and to support of cancer stem cells (130, 133, 136-138).

The aim of this study was to investigate PDGF-D and PDGFR β signaling in a heterogeneous tumor. For this purpose, we bred the *Pdgfd*^{-/-} animals with the RIP1-TAg2 mouse model of pancreatic neuroendocrine tumors (PanNET). The RIP1-TAg2 mice are engineered to express the oncogenic SV40 T antigen under the rat insulin promotor in the pancreatic islets, leading to hyperproliferation of islets and tumorigenesis. We found that PDGF-D deficiency delayed tumor growth by reducing cell proliferation, and thereby prolonging survival.

RIP1-TAg2 tumor formation has been studied previously in mice lacking the PDGF-B retention motif (*Pdgfb*^{ret/ret}) (168). In these mice, pericyte recruitment is severely disturbed in both normal and tumor vasculature, and has also been connected to an increased rate of metastasis. In consistence with our findings in Paper I, the expression of PDGF-D appeared to be mainly vascular in RIP1-TAg2; *Pdgfd*^{+/-} tumors. While vascular insufficiency and low presence of cytokine producing cells are common reasons for tumor growth stagnation, no effects were seen on angiogenesis, pericyte recruitment or immune cell infiltration. Also, the lack of PDGF-D did not affect the invasive nor metastatic properties of RIP1-TAg2 tumors. These findings are in coherence with our findings in Paper I, that the *Pdgfd*^{-/-} mice does not display any gross vascular abnormalities. Moreover, the metastatic properties are tightly connected to vascular integrity, and as the RIP1-TAg2; *Pdgfd*^{+/-} tumor vessel morphology was unaltered, the lack of alterations in metastatic frequency was not surprising.

To better understand the delayed growth in PDGF-D deficient tumors, we did an expression profiling by qPCR that showed an upregulation of *Pdgfb* in RIP1-TAg2; *Pdgfd*^{-/-} mice, compared to RIP1-TAg2; *Pdgfd*^{+/+} controls. This upregulation was likely aimed to make up for the lack PDGF-D, but clearly not enough to compensate for the reduced tumor growth, thus indicating a special role for PDGF-D in tumor development. Therefore, we explored the possibility of PDGF-D-responsive cells outside of the vascular compartment. By co-stainings of insulin and PDGFR β , in RIP1-TAg2; *Pdgfd*^{+/+} tumors, and fluorescence-activated cell sorting (FACS) analysis, we were able to identify a small subset of non-vascular tumor cells that were strongly expressing PDGFR β , and thus responsive to PDGF-D. Consistently, a rare PDGFR β ⁺ subset of cells was also present in the malignant β TC3 cell line, derived from RIP1-TAg2 tumors. This was a novel and highly interesting finding, as PDGFR β expression has been thought to be restricted to mural cells in the RIP1-TAg2 model (169). Interestingly, the β TC3 cells increased proliferation and formed a larger number of tumor spheroids upon treatment with PDGF-D, compared to PDGF-B, indicating a mechanistic difference between

PDGF-D and PDGF-B signaling. It is possible that the newly discovered PDGF-D co-receptor, NRP1, is involved in signaling modulation through altering, enhancing or prolonging signaling response. The function of NRP1 in modulation of PDGF-D signaling is addressed in Paper III.

To investigate the tumorigenic properties of PDGFR β ⁺ β TC3 cells, FACS sorted cells that were either PDGFR β ⁺ or PDGFR β ⁻ were subcutaneously transplanted into immunodeficient mice, and yielded tumors that were histologically indistinguishable from each other. Another FACS analysis showed that tumors formed from PDGFR β ⁻ cells had the same proportion of PDGFR β expression as tumors formed from PDGFR β ⁺ cells, thus indicating an inter-conversion between these two subsets of cells. We also verified that a subpopulation of PDGFR β ⁺ malignant cells is present also in human PanNET.

Taken together, we show that PDGF-D may contribute to PanNET tumor growth through angiocrine signaling, reinforcing functional malignant cell heterogeneity by stimulation of a rare PDGFR β ⁺ subset of tumor cells, which then engage in paracrine crosstalk with neighboring PDGFR β ⁻ cells to drive tumor progression. We also show that PDGFR β ⁻ tumor cells can switch into a PDGFR β ⁺ phenotype, to maintain a constant level of PDGF-D-responsive cells, thereby securing local access to tumorigenic signals. It would be interesting to further characterize NRP1 expression and its putative role in regulation of PDGF-D signaling in the progression of PanNET. Lastly, the notion that PDGFR β ⁺ malignant PanNET cells exist will likely impact the treatment regimens used for this type of cancer in the clinic. Sunitinib and other PDGFR β -targeting drugs might not only affect the vascular function, but also this rare population of highly tumorigenic cells, thus positioning these drugs as even more potent treatment options for PanNET than previously thought.

3.3 PAPER III – NEUROPILIN 1 IS A CO-RECEPTOR FOR PLATELET-DERIVED GROWTH FACTOR (PDGF)-D/PDGF RECEPTOR (PDGFR) β SIGNALING

The need for PDGF-B signaling through PDGFR β signaling in angiogenesis and vascular homeostasis is well documented. However, the role of PDGF-D is less clear. The existence of PDGF-D as a PDGFR β specific ligand, together with its need for extracellular activation, and the pathological implications from overexpression studies confirm that, once activated, PDGF-D is very potent in altering ECM homeostasis and other processes. Indeed, PDGF-D can induce responses separate from those induced by PDGF-B. As shown in Paper II, PDGF-D, but not PDGF-B, induces signaling in rare PDGFR β -expressing tumor cells, and contributes to tumor progression in PanNET in a manner that cannot be explained by differential activation. This could, however, be explained by modulation of signaling by a co-receptor.

There is a growing body of literature showing that NRP1 can modulate the signaling of both PDGFRs (42, 43, 140, 142, 143), but it does not bind to PDGFR β directly. Also, the affinity of PDGF-B binding to non-glycosylated NRP1 appears to be low (42, 143). Hence, this opens up for an additional, co-receptor-modulated role of PDGF-D as the other ligand for PDGFR β .

In this study, we discovered that PDGF-D shares some structural features with NRP1 binding VEGF ligands. In similarity with the NRP1-binding VEGF-A isoforms, VEGF-C, and VEGF-D, the PDGF-D growth factor domain has a conserved cysteine knot structure, accompanied by a C-terminal positively charged arginine residue, which in the VEGF ligands is crucial for NRP1 binding. However, these structural features are not seen in PDGF-B.

We show that PDGF-D specifically binds to NRP1 and induces PDGFR β /NRP1 complex formation in cells, both in *cis*- and in *trans*-conformation, and that binding leads to phosphorylation of PDGFR β , thus inducing downstream signaling. We also confirm that NRP1 binding of PDGF-B is only limited. Moreover, upon PDGF-D-stimulation of cultured cells, PDGF-D, NRP1 and PDGFR β were co-clustered and internalized together, suggesting that NRP1 alters the trafficking of PDGFR β . A similar mechanism has previously been shown for NRP1 in modulation of VEGF-signaling (170).

To our knowledge, this is the first study to assess the role of the novel PDGFs (PDGF-C or PDGF-D) as targets for NRP1 modulation. It makes sense that PDGF-D would be the preferred PDGFR β ligand for NRP modulation, considering that PDGF-D expression and activation are not coupled. With these new insights into the co-receptor function of NRP1 for PDGF-D/PDGFR β signaling, our study provides a mechanistic basis for differential signaling between PDGF-D and PDGF-B. It also adds complexity to the current hypotheses on NRP1 modulation of PDGF signaling and might even challenge some conclusions drawn from the previous reports.

Altogether, these findings will lead to a better understanding of the role(s) of PDGF-D, and thereby to improved development of tailored therapeutics for conditions where vascular function is disrupted, such as atherosclerosis and cancer.

4 FUTURE PERSPECTIVES

In the work included in this thesis, we present the *Pdgfd*^{-/-} mouse strain, and thereby the “PDGF family of knockouts” is finally complete. The mild phenotype of the *Pdgfd*^{-/-} mouse strain (described in Paper I) was not surprising to us, given the similarity between the PDGF-B and PDGFRβ phenotypes, indicating a certain amount of redundancy between PDGF-D and PDGF-B during physiological conditions. However, PDGF-B and PDGF-D signaling mechanisms differ in that PDGF-B is secreted in its active form, and retained close to the cell of origin through ECM binding, while PDGF-D is released in an inactive, diffusible form. Thus, PDGF-D expression and activation are uncoupled, and require the presence of additional activating factors, indicating a potent inducible function of PDGF-D, which might be important under specific conditions. Moreover, the identification of NRP1 as a co-receptor for PDGF-D, and its ability to form signaling complexes with PDGFRβ also in *trans* indicates that NRP1 does not only modulate signaling in the PDGFRβ -expressing cell, but also enables and potentiates cell-cell communication via direct adhesion to neighbouring NRP1-expressing cells. In my mind, this implies that a “main” function of PDGF-D is yet to be found.

In the vasculature, NRP1 is expressed by endothelial cells, and PDGFRβ by mural cells, thus the proposed signaling in *trans* is highly relevant for the communication between these two cell types. Interestingly, PDGF-D is commonly expressed around vascular bifurcations, and we have hypothesized that it is involved in regulation of local blood flow. Could it be that endothelial NRP1 acts in complex with mural PDGFRβ to regulate vascular contraction? This could be studied through an aortic ring assay, with and without PDGF-D stimulation. Another idea is that extra stability is needed in these areas due to turbulent blood flow, and that this is mediated through the NRP1-PDGF-D-PDGFRβ *trans* complexes.

It is also possible that PDGF-D has a role in angiocrine signaling with effects on cells in the tissues surrounding the vessels that could be somewhat similar to the mechanism described in Paper II. In this study, we present evidence that vascular PDGF-D reinforces the proliferative paracrine signaling of a rare population of PDGFRβ-expressing tumor cells to neighbouring tumor cells. Thus, the main site of action for PDGF-D might not be the vasculature, but some other tissue at some distance from the vessel. Interestingly, PDGFRβ expression has been reported in non-vascular progenitor cell populations and PDGF-D has been connected to stem cell signaling also in non-malignant conditions (60, 73, 74). Our reporter gene study indicated prominent expression of PDGF-D in several different epithelial tissues. Whereas epithelial cells are known to produce the PDGFRα-specific ligand PDGF-A (reviewed in (2)), the role of PDGFRβ in epithelial cells is not clear. However, the strong reporter gene expression of the PDGFRβ-specific PDGF-D is indicative of a role also for PDGFRβ signaling in the epithelium, possibly in a long-range paracrine manner similar to angiocrine signaling.




Figure 3. Possible physiological signaling modes for PDGF-D. PDGF-D is expressed by endothelial cells and pericytes, and activated by uPA, matriptase or possibly other unknown serine proteases. uPA is expressed by endothelial cells. (I) Short range paracrine or autocrine signaling in the vasculature. Pericytes express PDGFR β , and PDGF-D can either bind its receptor directly or engage the co-receptor NRP1 to form a complex. Complex formation can occur both in *cis* and in *trans*. (II) Angiocrine signaling, from the vasculature to cells in the surrounding tissue. The latent PDGF-D is diffusible and therefore not retained in the vasculature. Hypothetically, rare cells in the tissue express PDGFR β , and the same cells, or surrounding cells express activating serine proteases and the co-receptor NRP1. Thus, NRP1-modulation of PDGF-D/PDGFR β signaling in *cis* or *trans* could occur.

Now, the next step is to challenge the *Pdgfd*^{-/-} mice, to see if it has any phenotypic deviations compared to the wildtype mice. We have several projects involving challenges going on in the lab, and Paper II is derived from one of them. Another condition where PDGF-D is strongly implicated is atherosclerosis, and the SNPs in PDGF-D associated with coronary artery disease indicate that dysregulation of PDGF-D aggravates the disease. In an ongoing study, we have crossed the *Pdgfd*^{-/-} mice with the athero-prone *ApoE*^{-/-} mice. It will be very interesting to analyze the aortas from these mice to see if the plaques are differently constituted, and what cell populations that are altered.

In any case, it is possible that NRP1 holds the key to the function of PDGF-D, and this will be considered in future studies.

5 ACKNOWLEDGEMENTS

The process leading to the writing of this little book has been a long and journey full of sidetracks and dead ends. At times, searching for a phenotype in a seemingly healthy mouse feels like looking for a needle in a haystack. In spite of this, you are now finally holding my thesis in your hand. I would not have made it without you guys!

First of all, a big ‘Thank You’ to my main supervisor **Ulf**, for taking me in and for leading our group in such a good way. **Erika**, my co-supervisor and co-pilot on the PDGF-D project - thank you for believing in me and for always being there! You kept me on track, both when I got over-excited and when I felt like giving up. Also, thank you for all the fun we have had, and for advice on everything from science to house purchases and renovations, baby care, cooking and on-line shopping! ;) Also, big thanks to **Lars** who has been like a mini-co-supervisor on the NRP1 project lately. Thank you for introducing me to the world of protein biochemistry.

I also want to thank a few people who lead me to the UE lab: **Lotta**, former member of the KP lab - you were my "way in" as a totally green summer student! Thank you so much for referring me to the UE lab! **Carolina** and **Monika**, you showed me two extremely different ways to approach science, during that first summer at the Ludwig Institute. (Holy cow, those were two schizophrenic months...) Thank you guys for a really nice introduction to lab life! **Hong**, you were my supervisor when I came back for my master thesis - thank you for teaching me about cell culture!

Annika and **Eliane**, my fellow **Little Girls** - we started this journey together - and now, we finish our PhD era, just a few months apart! We shared *everything* during the first years, before Eliane moved to Lund and Annika and I had kids. I miss our endless dinners, all the travelling and shopping, sessions at Friskis and our deep sauna talks... Annika, your obsession with coffee and sweets has made a significant mark on my time at MBB. I want to thank you and the rest of **Team AF** for all of the nice *fika-fika* times and all of the treats that are always to be found around you! **Annelie**, you are always so cheery and keeping a firm hand on the steering wheel. (No, wait... I must be thinking of someone else...) **Sofia**, for being such a sweet and down-to-earth person. You really add some balance to this team. **Isolde** and **Karin** (and Annika again), I am still amazed that we all managed to become mothers in such a short time. Thanks for all the baby talk, and for sharing tips and tricks!

Philip, always with a smile almost as big as your hair! Thank you for interesting discussions about science and life in general. I am still sad that you, Erika and I never had the time to do the hiking-conference that we planned. **Anna**, I admire you for taking the decision to become a mum on your own, and I hope you will be back to your beloved yoga soon. Let me know when you start teaching classes!!! **Sebastian**, my lab bench neighbor! Thank you for all our nice chats about science and food. **Ingrid**, thank you for your pro tips on histology - your protocols are always the best! **Manuel** and **Milena**, also two histo stars and always helpful and friendly. **Christine**, good luck on your next missions, come what may!!!

Daniel and **Lars J**, you both do supercool science, and it has been a pleasure to have you and your growing groups in the same corridor! **Linda** - thank you for our discussions on the subway home, and for your new angles and your support with the PDGF-D project. **Josefin**, I know you will be excellent as an MSL! **Yixin**, my back-to-back office mate, I think we have been a great team when it comes to answering each other's stupid questions and problems! **Aga** - good luck with your little one!

Also, a big thanks to the rest of the people in Vascular Biology and in Matrix, for good times and for sharing knowledge and reagents. **Frank, Joanna, Mirela, Yi, Michael, Christina, Guillem, Tian, Jong-Wook, Arancha, Hassan, Hosein, Anne-May, Patricia, Juha** and **Lwaki** - it has been a pleasure!

Friends in groups that have moved from MBB: **KP lab** - **Kristian**, thank you for great collaboration! Your group has really developed since I came to the lab, and you always manage find the nicest people: **Matteo** - our Italian drama king, with the warmest heart! Madre mía, I miss you up here!!! **Nik** and **Micha** - I'm so glad that I got to know you before you moved to Lund, and I also miss **Pernilla, Sara** and **Marcela**, the tea-loving, super sweet KP girls! The **CBZ** gang - **Jen**, you were my office mate for a long time, thanks for all your input on science and for your friendship! **Elizabeth, Barbara, Maarja, Maya, Johanna, Konstantin** and **Colin** thank you for all the nice chats and parties! And **Fuxe** group - one day you were here and the next you were not... Anyway, thank you **Jonas, Azi, Ana, Joel, Meifong, Vedrana, Tove** and **Jill** for sharing antibodies and good times!

Claudia - con tigo en el pasillo, todo está bien! But when you are not there, we notice right away that things are not running as smoothly - thank you for making lab life so much easier. Last but not least, **Gizella** - wow, what a difference you have made in the corridor! Thank you for all the "fixing", and for bringing your warmth and energy in everything you do - you are such a star!!!

I also want to thank two supportive husbands, (Annika's) **Martin** and (Sofia's) **Björn**, for being part of the vivid lab-life before all the babies started arriving. Thank you for all the nice dinners and parties!

Även om det inte verkat så sista tiden så har jag ju ett liv utanför labbet också...

Lisa, du är den som jag känt längst av alla - tänk vad mycket vi varit med om sedan vi träffades första gången i den där snödrivan! Jag är så glad att jag har dig i mitt liv, och det är alltid lika trevligt att hänga med dig och **Martin**.

Jemima, min mammagrupsbestie! Vi klickade direkt och kom varandra väldigt nära på kort tid, jag är så glad att vi lärt känna varandra och att du hade den goda smaken att föda barn samtidigt som jag. ;) Tack för allt trevligt häng med dig, **Henrik** och lilla **Arvid** i Fruängen, och hoppas det fortsätter framöver, trots flytt!

Skog och **Miranda**, tack för alla spontana vardagsmiddagar och skratt, snaaart ska vi börja

bjuda igen för det senaste halvåret...

Henning, vi har haft många djupa diskussioner om allt mellan himmel och jord! Kom tillbaka till Stockholm snart, jag saknar dig!

Finaste **Fanny**, min bästis från Biomedicinprogrammet - tack för att du finns och för att du är en sådan inspirationskälla! Älskar att du och din **Jonas** dessutom delar vår passion för sällskapsspel. Resten av biomedicinbrudarna, **Anna**, **Sofia W**, **Sofia A**, **Mariann** och **Niosha**, jag är så glad att vi fortsatt med våra månatliga middagar - vi har verkligen blivit ett sammansvetsat litet gäng, love you!

Mamma och **pappa**, tack för att ni alltid uppmuntrat och stöttat mig och mina bröder att göra det vi gillar! **Kristofer**, tänk att du hann publicera en vetenskaplig artikel före mig trots att du inte ens blev doktorand. Så kul att ha en till forskare i familjen! Vill även tacka dig, **Jonas** och **Rasmus**, för allt härligt syskonhång! Alltid lika trevligt när vi är hemma alla samtidigt och särskilt när **Sara**, **Linnéa** och **Elin** är med.

Mormor och **morfar**, för att ni stöttat, curlat och skämt bort oss syskon när vi växte upp. Morfar, jag hade aldrig blivit den jag är idag om det inte vore för dig - du inspirerade mig att läsa vidare, och fick mig att intressera mig för teknik och naturvetenskap. **Farmor**, jag älskar att du kör ditt eget race, med resor och annat. Jag vill bli mer som du! Moster **Annika**, tack för alla pep-talks. Din drivkraft och din smittande entusiasm är fantastisk!

Familjen Selling: **Maud** och **Glenn**, tack för att ni tagit emot mig så öppenhjärtigt i familjen, bättre svärföräldrar kan man inte önska sig. **Malin**, tack för snabbkursen i Illustrator - du anar inte vad mycket lättare du gjorde det för mig, nu på slutspurt! Tack även för allt skönt häng med dig, **Anton** och **Mattias**!

Slutligen, mina fina grabbar. **Felix**, älskade lilla skrutt, tack för att du lärt mig vad som är viktigast här i livet, och för att du tvingar mig att koppla bort jobbet! Fina, fina **Jimmy**, tack för att du finns. Jag vet inte vad jag skulle göra utan dig, och jag hade aldrig tagit mig igenom det här utan ditt stöd. Jag älskar dig!

6 REFERENCES

1. Heldin CH, Westermark B. Mechanism of action and in vivo role of platelet-derived growth factor. *Physiological reviews*. 1999;79(4):1283-316. Epub 1999/10/03.
2. Andrae J, Gallini R, Betsholtz C. Role of platelet-derived growth factors in physiology and medicine. *Genes & development*. 2008;22(10):1276-312. Epub 2008/05/17.
3. Fredriksson L, Li H, Eriksson U. The PDGF family: four gene products form five dimeric isoforms. *Cytokine & growth factor reviews*. 2004;15(4):197-204. Epub 2004/06/23.
4. Bork P, Beckmann G. The CUB domain. A widespread module in developmentally regulated proteins. *Journal of molecular biology*. 1993;231(2):539-45. Epub 1993/05/20.
5. Bergsten E, Uutela M, Li X, Pietras K, Ostman A, Heldin CH, et al. PDGF-D is a specific, protease-activated ligand for the PDGF beta-receptor. *Nature cell biology*. 2001;3(5):512-6. Epub 2001/05/02.
6. LaRochelle WJ, Jeffers M, McDonald WF, Chillakuru RA, Giese NA, Lokker NA, et al. PDGF-D, a new protease-activated growth factor. *Nature cell biology*. 2001;3(5):517-21. Epub 2001/05/02.
7. Eppig JT, Blake JA, Bult CJ, Kadin JA, Richardson JE. The Mouse Genome Database (MGD): facilitating mouse as a model for human biology and disease. *Nucleic acids research*. 2015;43(Database issue):D726-36. Epub 2014/10/29.
8. Uutela M, Lauren J, Bergsten E, Li X, Horelli-Kuitunen N, Eriksson U, et al. Chromosomal location, exon structure, and vascular expression patterns of the human PDGFC and PDGFD genes. *Circulation*. 2001;103(18):2242-7. Epub 2001/05/23.
9. Ustach CV, Kim HR. Platelet-derived growth factor D is activated by urokinase plasminogen activator in prostate carcinoma cells. *Molecular and cellular biology*. 2005;25(14):6279-88. Epub 2005/07/01.
10. Ehnman M, Li H, Fredriksson L, Pietras K, Eriksson U. The uPA/uPAR system regulates the bioavailability of PDGF-DD: implications for tumour growth. *Oncogene*. 2009;28(4):534-44. Epub 2008/11/11.
11. Ustach CV, Huang W, Conley-LaComb MK, Lin CY, Che M, Abrams J, et al. A novel signaling axis of matriptase/PDGF-D/ss-PDGFR in human prostate cancer. *Cancer research*. 2010;70(23):9631-40. Epub 2010/11/26.
12. Rorsman F, Bywater M, Knott TJ, Scott J, Betsholtz C. Structural characterization of the human platelet-derived growth factor A-chain cDNA and gene: alternative exon usage predicts two different precursor proteins. *Molecular and cellular biology*. 1988;8(2):571-7. Epub 1988/02/01.
13. Ostman A, Andersson M, Betsholtz C, Westermark B, Heldin CH. Identification of a cell retention signal in the B-chain of platelet-derived growth factor and in the long splice version of the A-chain. *Cell regulation*. 1991;2(7):503-12. Epub 1991/07/01.
14. Raines EW, Ross R. Compartmentalization of PDGF on extracellular binding sites dependent on exon-6-encoded sequences. *The Journal of cell biology*. 1992;116(2):533-43. Epub 1992/01/01.
15. Andrae J, Ehrencrona H, Gallini R, Lal M, Ding H, Betsholtz C. Analysis of mice lacking the heparin-binding splice isoform of platelet-derived growth factor A. *Molecular and cellular biology*. 2013;33(20):4030-40. Epub 2013/08/14.
16. Zhuo Y, Hoyle GW, Zhang J, Morris G, Lasky JA. A novel murine PDGF-D splicing variant results in significant differences in peptide expression and function. *Biochemical and biophysical research communications*. 2003;308(1):126-32. Epub 2003/08/02.
17. Ehnman M, Missiaglia E, Folestad E, Selfe J, Strell C, Thway K, et al. Distinct effects of ligand-induced PDGFRalpha and PDGFRbeta signaling in the human rhabdomyosarcoma tumor cell and stroma cell compartments. *Cancer research*. 2013;73(7):2139-49. Epub 2013/01/23.
18. Heldin CH. Targeting the PDGF signaling pathway in tumor treatment. *Cell communication and signaling : CCS*. 2013;11:97. Epub 2013/12/24.
19. Klinghoffer RA, Mueting-Nelsen PF, Faerman A, Shani M, Soriano P. The two PDGF receptors maintain conserved signaling in vivo despite divergent embryological functions. *Molecular cell*. 2001;7(2):343-54. Epub 2001/03/10.
20. Goh LK, Sorkin A. Endocytosis of receptor tyrosine kinases. *Cold Spring Harbor perspectives in biology*. 2013;5(5):a017459. Epub 2013/05/03.
21. Orr-Urtreger A, Bedford MT, Do MS, Eisenbach L, Lonai P. Developmental expression of the alpha receptor for platelet-derived growth factor, which is deleted in the embryonic lethal Patch mutation. *Development*. 1992;115(1):289-303. Epub 1992/05/01.

22. Ataliotis P, Mercola M. Distribution and functions of platelet-derived growth factors and their receptors during embryogenesis. *International review of cytology*. 1997;172:95-127. Epub 1997/01/01.
23. Lindahl P, Johansson BR, Leveen P, Betsholtz C. Pericyte loss and microaneurysm formation in PDGF-B-deficient mice. *Science*. 1997;277(5323):242-5. Epub 1997/07/11.
24. Raines EW. PDGF and cardiovascular disease. *Cytokine & growth factor reviews*. 2004;15(4):237-54. Epub 2004/06/23.
25. Hamada T, Ui-Tei K, Imaki J, Miyata Y. Molecular cloning of SCDGF-B, a novel growth factor homologous to SCDGF/PDGF-C/fallotin. *Biochemical and biophysical research communications*. 2001;280(3):733-7. Epub 2001/02/13.
26. Gladh H, Folestad EB, Muhl L, Ehnman M, Tannenbergs P, Lawrence AL, et al. Mice Lacking Platelet-Derived Growth Factor D Display a Mild Vascular Phenotype. *PloS one*. 2016;11(3):e0152276. Epub 2016/04/01.
27. Chen J, Han Y, Lin C, Zhen Y, Song X, Teng S, et al. PDGF-D contributes to neointimal hyperplasia in rat model of vessel injury. *Biochemical and biophysical research communications*. 2005;329(3):976-83. Epub 2005/03/09.
28. Karvinen H, Rutanen J, Leppanen O, Lach R, Levonen AL, Eriksson U, et al. PDGF-C and -D and their receptors PDGFR-alpha and PDGFR-beta in atherosclerotic human arteries. *European journal of clinical investigation*. 2009;39(4):320-7. Epub 2009/03/19.
29. Ponten A, Folestad EB, Pietras K, Eriksson U. Platelet-derived growth factor D induces cardiac fibrosis and proliferation of vascular smooth muscle cells in heart-specific transgenic mice. *Circulation research*. 2005;97(10):1036-45. Epub 2005/10/15.
30. Van Den Akker NM, Lie-Venema H, Maas S, Eralp I, DeRuiter MC, Poelmann RE, et al. Platelet-derived growth factors in the developing avian heart and maturing coronary vasculature. *Developmental dynamics : an official publication of the American Association of Anatomists*. 2005;233(4):1579-88. Epub 2005/06/24.
31. Changsirikulchai S, Hudkins KL, Goodpaster TA, Volpone J, Topouzis S, Gilbertson DG, et al. Localization of platelet derived growth factor-D (PDGF-D) and its receptor in human kidneys. *J Am Soc Nephrol*. 2002;13:78A-A.
32. Buhl EM, Djudjaj S, Babickova J, Klinkhammer BM, Folestad E, Borkham-Kamphorst E, et al. The role of PDGF-D in healthy and fibrotic kidneys. *Kidney international*. 2016;89(4):848-61. Epub 2016/03/01.
33. Ray S, Gao C, Wyatt K, Fariss RN, Bundek A, Zelenka P, et al. Platelet-derived growth factor D, tissue-specific expression in the eye, and a key role in control of lens epithelial cell proliferation. *The Journal of biological chemistry*. 2005;280(9):8494-502. Epub 2004/12/22.
34. Bozanic D, Bocina I, Saraga-Babic M. Involvement of cytoskeletal proteins and growth factor receptors during development of the human eye. *Anatomy and embryology*. 2006;211(5):367-77. Epub 2006/04/06.
35. Bennett JA, Singh KP, Unnisa Z, Welle SL, Gasiewicz TA. Deficiency in Aryl Hydrocarbon Receptor (AHR) Expression throughout Aging Alters Gene Expression Profiles in Murine Long-Term Hematopoietic Stem Cells. *PloS one*. 2015;10(7):e0133791. Epub 2015/07/25.
36. Kim WS, Park HS, Sung JH. The pivotal role of PDGF and its receptor isoforms in adipose-derived stem cells. *Histology and histopathology*. 2015;30(7):793-9. Epub 2015/02/26.
37. Hamada T, Ui-Tei K, Imaki J, Takahashi F, Onodera H, Mishima T, et al. The expression of SCDGF/PDGF-C/fallotin and SCDGF-B/PDGF-D in the rat central nervous system. *Mechanisms of development*. 2002;112(1-2):161-4. Epub 2002/02/19.
38. Fredriksson L, Li H, Fieber C, Li X, Eriksson U. Tissue plasminogen activator is a potent activator of PDGF-CC. *The EMBO journal*. 2004;23(19):3793-802. Epub 2004/09/17.
39. Rogers KW, Schier AF. Morphogen gradients: from generation to interpretation. *Annual review of cell and developmental biology*. 2011;27:377-407. Epub 2011/08/02.
40. Loukinova E, Ranganathan S, Kuznetsov S, Gorlatova N, Migliorini MM, Loukinov D, et al. Platelet-derived growth factor (PDGF)-induced tyrosine phosphorylation of the low density lipoprotein receptor-related protein (LRP). Evidence for integrated co-receptor function between LRP and the PDGF. *The Journal of biological chemistry*. 2002;277(18):15499-506. Epub 2002/02/21.
41. Kiyan J, Kiyan R, Haller H, Dumler I. Urokinase-induced signaling in human vascular smooth muscle cells is mediated by PDGFR-beta. *The EMBO journal*. 2005;24(10):1787-97. Epub 2005/05/13.

42. Ball SG, Bayley C, Shuttleworth CA, Kielty CM. Neuropilin-1 regulates platelet-derived growth factor receptor signalling in mesenchymal stem cells. *The Biochemical journal*. 2010;427(1):29-40. Epub 2010/01/28.
43. Banerjee S, Sengupta K, Dhar K, Mehta S, D'Amore PA, Dhar G, et al. Breast cancer cells secreted platelet-derived growth factor-induced motility of vascular smooth muscle cells is mediated through neuropilin-1. *Molecular carcinogenesis*. 2006;45(11):871-80. Epub 2006/07/19.
44. Takayama Y, May P, Anderson RG, Herz J. Low density lipoprotein receptor-related protein 1 (LRP1) controls endocytosis and c-CBL-mediated ubiquitination of the platelet-derived growth factor receptor beta (PDGFR beta). *The Journal of biological chemistry*. 2005;280(18):18504-10. Epub 2005/03/09.
45. Nakayama A, Nakayama M, Turner CJ, Hoing S, Lepore JJ, Adams RH. Ephrin-B2 controls PDGFRbeta internalization and signaling. *Genes & development*. 2013;27(23):2576-89. Epub 2013/12/04.
46. Mendelson K, Swendeman S, Saftig P, Blobel CP. Stimulation of platelet-derived growth factor receptor beta (PDGFRbeta) activates ADAM17 and promotes metalloproteinase-dependent cross-talk between the PDGFRbeta and epidermal growth factor receptor (EGFR) signaling pathways. *The Journal of biological chemistry*. 2010;285(32):25024-32. Epub 2010/06/10.
47. Lei H, Kazlauskas A. Growth factors outside of the platelet-derived growth factor (PDGF) family employ reactive oxygen species/Src family kinases to activate PDGF receptor alpha and thereby promote proliferation and survival of cells. *The Journal of biological chemistry*. 2009;284(10):6329-36. Epub 2009/01/08.
48. Shimizu H, Hirose Y, Nishijima F, Tsubakihara Y, Miyazaki H. ROS and PDGF-beta [corrected] receptors are critically involved in indoxyl sulfate actions that promote vascular smooth muscle cell proliferation and migration. *American journal of physiology Cell physiology*. 2009;297(2):C389-96. Epub 2009/06/06.
49. Tallquist M, Kazlauskas A. PDGF signaling in cells and mice. *Cytokine & growth factor reviews*. 2004;15(4):205-13. Epub 2004/06/23.
50. Ding H, Wu X, Bostrom H, Kim I, Wong N, Tsoi B, et al. A specific requirement for PDGF-C in palate formation and PDGFR-alpha signaling. *Nature genetics*. 2004;36(10):1111-6. Epub 2004/09/14.
51. Fredriksson L, Nilsson I, Su EJ, Andrae J, Ding H, Betsholtz C, et al. Platelet-derived growth factor C deficiency in C57BL/6 mice leads to abnormal cerebral vascularization, loss of neuroependymal integrity, and ventricular abnormalities. *The American journal of pathology*. 2012;180(3):1136-44. Epub 2012/01/11.
52. Leveen P, Pekny M, Gebre-Medhin S, Swolin B, Larsson E, Betsholtz C. Mice deficient for PDGF B show renal, cardiovascular, and hematological abnormalities. *Genes & development*. 1994;8(16):1875-87. Epub 1994/08/15.
53. Soriano P. Abnormal kidney development and hematological disorders in PDGF beta-receptor mutant mice. *Genes & development*. 1994;8(16):1888-96. Epub 1994/08/15.
54. Hellstrom M, Kalen M, Lindahl P, Abramsson A, Betsholtz C. Role of PDGF-B and PDGFR-beta in recruitment of vascular smooth muscle cells and pericytes during embryonic blood vessel formation in the mouse. *Development*. 1999;126(14):3047-55. Epub 1999/06/22.
55. Jakobsson L, Franco CA, Bentley K, Collins RT, Ponsioen B, Aspalter IM, et al. Endothelial cells dynamically compete for the tip cell position during angiogenic sprouting. *Nature cell biology*. 2010;12(10):943-53. Epub 2010/09/28.
56. Hirschi KK, Rohovsky SA, Beck LH, Smith SR, D'Amore PA. Endothelial cells modulate the proliferation of mural cell precursors via platelet-derived growth factor-BB and heterotypic cell contact. *Circulation research*. 1999;84(3):298-305. Epub 1999/02/19.
57. Bjarnegard M, Enge M, Norlin J, Gustafsdottir S, Fredriksson S, Abramsson A, et al. Endothelium-specific ablation of PDGFB leads to pericyte loss and glomerular, cardiac and placental abnormalities. *Development*. 2004;131(8):1847-57. Epub 2004/04/16.
58. Mazzone M, Dettori D, Leite de Oliveira R, Loges S, Schmidt T, Jonckx B, et al. Heterozygous deficiency of PHD2 restores tumor oxygenation and inhibits metastasis via endothelial normalization. *Cell*. 2009;136(5):839-51. Epub 2009/02/17.
59. Hellstrom M, Gerhardt H, Kalen M, Li X, Eriksson U, Wolburg H, et al. Lack of pericytes leads to endothelial hyperplasia and abnormal vascular morphogenesis. *J Cell Biol*. 2001;153(3):543-53. Epub 2001/05/02.
60. Basciani S, De Luca G, Dolci S, Brama M, Arizzi M, Mariani S, et al. Platelet-derived growth factor receptor beta-subtype regulates proliferation and migration of gonocytes. *Endocrinology*. 2008;149(12):6226-35. Epub 2008/08/09.
61. Homsy J, Daud AI. Spectrum of activity and mechanism of action of VEGF/PDGF inhibitors. *Cancer control : journal of the Moffitt Cancer Center*. 2007;14(3):285-94. Epub 2007/07/07.

62. Heuchel R, Berg A, Tallquist M, Ahlen K, Reed RK, Rubin K, et al. Platelet-derived growth factor beta receptor regulates interstitial fluid homeostasis through phosphatidylinositol-3' kinase signaling. *Proceedings of the National Academy of Sciences of the United States of America*. 1999;96(20):11410-5. Epub 1999/09/29.
63. Hawthorne T, Giot L, Blake L, Kuang B, Gerwien R, Smithson G, et al. A phase I study of CR002, a fully-human monoclonal antibody against platelet-derived growth factor-D. *International journal of clinical pharmacology and therapeutics*. 2008;46(5):236-44. Epub 2008/06/10.
64. Lindblom P, Gerhardt H, Liebner S, Abramsson A, Enge M, Hellstrom M, et al. Endothelial PDGF-B retention is required for proper investment of pericytes in the microvessel wall. *Genes & development*. 2003;17(15):1835-40. Epub 2003/08/05.
65. Armulik A, Genove G, Mae M, Nisancioglu MH, Wallgard E, Niaudet C, et al. Pericytes regulate the blood-brain barrier. *Nature*. 2010;468(7323):557-61. Epub 2010/10/15.
66. Pierce GF, Mustoe TA, Altmann BW, Deuel TF, Thomason A. Role of platelet-derived growth factor in wound healing. *Journal of cellular biochemistry*. 1991;45(4):319-26. Epub 1991/04/01.
67. Fang L, Yan Y, Komuves LG, Yonkovich S, Sullivan CM, Stringer B, et al. PDGF C is a selective alpha platelet-derived growth factor receptor agonist that is highly expressed in platelet alpha granules and vascular smooth muscle. *Arteriosclerosis, thrombosis, and vascular biology*. 2004;24(4):787-92. Epub 2004/04/06.
68. Robson MC, Phillips LG, Thomason A, Altmann BW, Pence PC, Heggers JP, et al. Recombinant human platelet-derived growth factor-BB for the treatment of chronic pressure ulcers. *Annals of plastic surgery*. 1992;29(3):193-201. Epub 1992/09/01.
69. Uutela M, Wirzenius M, Paavonen K, Rajantie I, He Y, Karpanen T, et al. PDGF-D induces macrophage recruitment, increased interstitial pressure, and blood vessel maturation during angiogenesis. *Blood*. 2004;104(10):3198-204. Epub 2004/07/24.
70. Wagsater D, Zhu C, Bjorck HM, Eriksson P. Effects of PDGF-C and PDGF-D on monocyte migration and MMP-2 and MMP-9 expression. *Atherosclerosis*. 2009;202(2):415-23. Epub 2008/06/25.
71. Germain S, Monnot C, Muller L, Eichmann A. Hypoxia-driven angiogenesis: role of tip cells and extracellular matrix scaffolding. *Current opinion in hematology*. 2010;17(3):245-51. Epub 2010/03/24.
72. Rodt SA, Ahlen K, Berg A, Rubin K, Reed RK. A novel physiological function for platelet-derived growth factor-BB in rat dermis. *The Journal of physiology*. 1996;495 (Pt 1):193-200. Epub 1996/08/15.
73. Schwab KE, Gargett CE. Co-expression of two perivascular cell markers isolates mesenchymal stem-like cells from human endometrium. *Hum Reprod*. 2007;22(11):2903-11. Epub 2007/09/18.
74. Hye Kim J, Gyu Park S, Kim WK, Song SU, Sung JH. Functional regulation of adipose-derived stem cells by PDGF-D. *Stem Cells*. 2015;33(2):542-56. Epub 2014/10/22.
75. Isaka Y, Fujiwara Y, Ueda N, Kaneda Y, Kamada T, Imai E. Glomerulosclerosis induced by in vivo transfection of transforming growth factor-beta or platelet-derived growth factor gene into the rat kidney. *The Journal of clinical investigation*. 1993;92(6):2597-601. Epub 1993/12/01.
76. Zeisberg M, Kalluri R. Cellular mechanisms of tissue fibrosis. 1. Common and organ-specific mechanisms associated with tissue fibrosis. *American journal of physiology Cell physiology*. 2013;304(3):C216-25. Epub 2012/12/21.
77. Bonner JC. Regulation of PDGF and its receptors in fibrotic diseases. *Cytokine & growth factor reviews*. 2004;15(4):255-73. Epub 2004/06/23.
78. Zhao T, Zhao W, Chen Y, Li VS, Meng W, Sun Y. Platelet-derived growth factor-D promotes fibrogenesis of cardiac fibroblasts. *American journal of physiology Heart and circulatory physiology*. 2013;304(12):H1719-26. Epub 2013/04/16.
79. Charni Chaabane S, Coomans de Brachene A, Essaghir A, Velghe A, Lo Re S, Stockis J, et al. PDGF-D expression is down-regulated by TGFbeta in fibroblasts. *PloS one*. 2014;9(10):e108656. Epub 2014/10/04.
80. van Roeyen CR, Eitner F, Boor P, Moeller MJ, Raffetseder U, Hanssen L, et al. Induction of progressive glomerulonephritis by podocyte-specific overexpression of platelet-derived growth factor-D. *Kidney international*. 2011;80(12):1292-305. Epub 2011/08/26.
81. Breitkopf K, Roeyen C, Sawitza I, Wickert L, Floege J, Gressner AM. Expression patterns of PDGF-A, -B, -C and -D and the PDGF-receptors alpha and beta in activated rat hepatic stellate cells (HSC). *Cytokine*. 2005;31(5):349-57. Epub 2005/07/26.
82. Ostendorf T, Rong S, Boor P, Wiedemann S, Kunter U, Haubold U, et al. Antagonism of PDGF-D by human antibody CR002 prevents renal scarring in experimental glomerulonephritis. *J Am Soc Nephrol*. 2006;17(4):1054-62. Epub 2006/03/03.

83. Borkham-Kamphorst E, van Roeyen CR, Ostendorf T, Floege J, Gressner AM, Weiskirchen R. Pro-fibrogenic potential of PDGF-D in liver fibrosis. *Journal of hepatology*. 2007;46(6):1064-74. Epub 2007/04/03.
84. Kinnman N, Gorla O, Wendum D, Gendron MC, Rey C, Poupon R, et al. Hepatic stellate cell proliferation is an early platelet-derived growth factor-mediated cellular event in rat cholestatic liver injury. *Laboratory investigation; a journal of technical methods and pathology*. 2001;81(12):1709-16. Epub 2001/12/14.
85. Borkham-Kamphorst E, Meurer SK, Van de Leur E, Haas U, Tihaa L, Weiskirchen R. PDGF-D signaling in portal myofibroblasts and hepatic stellate cells proves identical to PDGF-B via both PDGF receptor type alpha and beta. *Cellular signalling*. 2015;27(7):1305-14. Epub 2015/03/31.
86. Ostendorf T, van Roeyen CR, Peterson JD, Kunter U, Eitner F, Hamad AJ, et al. A fully human monoclonal antibody (CR002) identifies PDGF-D as a novel mediator of mesangioproliferative glomerulonephritis. *J Am Soc Nephrol*. 2003;14(9):2237-47. Epub 2003/08/26.
87. Ahmad A, Wang Z, Kong D, Ali R, Ali S, Banerjee S, et al. Platelet-derived growth factor-D contributes to aggressiveness of breast cancer cells by up-regulating Notch and NF-kappaB signaling pathways. *Breast cancer research and treatment*. 2011;126(1):15-25. Epub 2010/04/10.
88. Kalluri R, Weinberg RA. The basics of epithelial-mesenchymal transition. *The Journal of clinical investigation*. 2009;119(6):1420-8. Epub 2009/06/03.
89. Boor P, Konieczny A, Villa L, Kunter U, van Roeyen CR, LaRochelle WJ, et al. PDGF-D inhibition by CR002 ameliorates tubulointerstitial fibrosis following experimental glomerulonephritis. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association*. 2007;22(5):1323-31. Epub 2007/02/20.
90. He C, Medley SC, Hu T, Hinsdale ME, Lupu F, Virmani R, et al. PDGFRbeta signalling regulates local inflammation and synergizes with hypercholesterolaemia to promote atherosclerosis. *Nature communications*. 2015;6:7770. Epub 2015/07/18.
91. Hansson GK. Inflammation, atherosclerosis, and coronary artery disease. *The New England journal of medicine*. 2005;352(16):1685-95. Epub 2005/04/22.
92. Doran AC, Meller N, McNamara CA. Role of smooth muscle cells in the initiation and early progression of atherosclerosis. *Arteriosclerosis, thrombosis, and vascular biology*. 2008;28(5):812-9. Epub 2008/02/16.
93. Thomas JA, Deaton RA, Hastings NE, Shang Y, Moehle CW, Eriksson U, et al. PDGF-DD, a novel mediator of smooth muscle cell phenotypic modulation, is upregulated in endothelial cells exposed to atherosclerosis-prone flow patterns. *American journal of physiology Heart and circulatory physiology*. 2009;296(2):H442-52. Epub 2008/11/26.
94. A genome-wide association study in Europeans and South Asians identifies five new loci for coronary artery disease. *Nature genetics*. 2011;43(4):339-44. Epub 2011/03/08.
95. Bai Y, Chen J, Sun K, Wang Y, Hui R. A functional variant in promoter region of platelet-derived growth factor-D is probably associated with intracerebral hemorrhage. *Journal of neuroinflammation*. 2012;9:26. Epub 2012/02/01.
96. Costello JP, Mohanakumar T, Nath DS. Mechanisms of chronic cardiac allograft rejection. *Texas Heart Institute journal / from the Texas Heart Institute of St Luke's Episcopal Hospital, Texas Children's Hospital*. 2013;40(4):395-9. Epub 2013/10/02.
97. Tuuminen R, Nykanen A, Keranen MA, Krebs R, Alitalo K, Koskinen PK, et al. The effect of platelet-derived growth factor ligands in rat cardiac allograft vasculopathy and fibrosis. *Transplantation proceedings*. 2006;38(10):3271-3. Epub 2006/12/19.
98. Su EJ, Fredriksson L, Geyer M, Folestad E, Cale J, Andrae J, et al. Activation of PDGF-CC by tissue plasminogen activator impairs blood-brain barrier integrity during ischemic stroke. *Nature medicine*. 2008;14(7):731-7. Epub 2008/06/24.
99. Su EJ, Fredriksson L, Schielke GP, Eriksson U, Lawrence DA. Tissue plasminogen activator-mediated PDGF signaling and neurovascular coupling in stroke. *Journal of thrombosis and haemostasis : JTH*. 2009;7 Suppl 1:155-8. Epub 2009/07/28.
100. Lewandowski SA, Nilsson I, Fredriksson L, Lonnerberg P, Muhl L, Zeitelhofer M, et al. Presymptomatic activation of the PDGF-CC pathway accelerates onset of ALS neurodegeneration. *Acta neuropathologica*. 2016;131(3):453-64. Epub 2015/12/22.
101. Hanahan D, Weinberg RA. The hallmarks of cancer. *Cell*. 2000;100(1):57-70. Epub 2000/01/27.
102. Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell*. 2011;144(5):646-74. Epub 2011/03/08.

103. Collins T, Ginsburg D, Boss JM, Orkin SH, Pober JS. Cultured human endothelial cells express platelet-derived growth factor B chain: cDNA cloning and structural analysis. *Nature*. 1985;316(6030):748-50. Epub 1985/08/22.
104. Doolittle RF, Hunkapiller MW, Hood LE, Devare SG, Robbins KC, Aaronson SA, et al. Simian sarcoma virus onc gene, v-sis, is derived from the gene (or genes) encoding a platelet-derived growth factor. *Science*. 1983;221(4607):275-7. Epub 1983/07/15.
105. Pietras K, Sjoblom T, Rubin K, Heldin CH, Ostman A. PDGF receptors as cancer drug targets. *Cancer cell*. 2003;3(5):439-43. Epub 2003/06/05.
106. Shimizu A, O'Brien KP, Sjoblom T, Pietras K, Buchdunger E, Collins VP, et al. The dermatofibrosarcoma protuberans-associated collagen type Ialpha1/platelet-derived growth factor (PDGF) B-chain fusion gene generates a transforming protein that is processed to functional PDGF-BB. *Cancer research*. 1999;59(15):3719-23. Epub 1999/08/14.
107. Cools J, DeAngelo DJ, Gotlib J, Stover EH, Legare RD, Cortes J, et al. A tyrosine kinase created by fusion of the PDGFRA and FIP1L1 genes as a therapeutic target of imatinib in idiopathic hypereosinophilic syndrome. *The New England journal of medicine*. 2003;348(13):1201-14. Epub 2003/03/28.
108. Golub TR, Barker GF, Lovett M, Gilliland DG. Fusion of PDGF receptor beta to a novel ets-like gene, tel, in chronic myelomonocytic leukemia with t(5;12) chromosomal translocation. *Cell*. 1994;77(2):307-16. Epub 1994/04/22.
109. Zheng S, Chen LR, Wang HJ, Chen SZ. Analysis of mutation and expression of c-kit and PDGFR-alpha gene in gastrointestinal stromal tumor. *Hepato-gastroenterology*. 2007;54(80):2285-90. Epub 2008/02/13.
110. Joensuu H, Pupa M, Sihto H, Tynnenen O, Nupponen NN. Amplification of genes encoding KIT, PDGFRalpha and VEGFR2 receptor tyrosine kinases is frequent in glioblastoma multiforme. *The Journal of pathology*. 2005;207(2):224-31. Epub 2005/07/16.
111. Demoulin JB, Essaghir A. PDGF receptor signaling networks in normal and cancer cells. *Cytokine & growth factor reviews*. 2014;25(3):273-83. Epub 2014/04/08.
112. Cortez E, Roswall P, Pietras K. Functional subsets of mesenchymal cell types in the tumor microenvironment. *Seminars in cancer biology*. 2014;25:3-9. Epub 2014/01/15.
113. Furuhashi M, Sjoblom T, Abramsson A, Ellingsen J, Micke P, Li H, et al. Platelet-derived growth factor production by B16 melanoma cells leads to increased pericyte abundance in tumors and an associated increase in tumor growth rate. *Cancer research*. 2004;64(8):2725-33. Epub 2004/04/17.
114. Franco M, Roswall P, Cortez E, Hanahan D, Pietras K. Pericytes promote endothelial cell survival through induction of autocrine VEGF-A signaling and Bcl-w expression. *Blood*. 2011;118(10):2906-17. Epub 2011/07/23.
115. Pietras K, Ostman A, Sjoquist M, Buchdunger E, Reed RK, Heldin CH, et al. Inhibition of platelet-derived growth factor receptors reduces interstitial hypertension and increases transcapillary transport in tumors. *Cancer research*. 2001;61(7):2929-34. Epub 2001/04/18.
116. Wang Z, Kong D, Li Y, Sarkar FH. PDGF-D signaling: a novel target in cancer therapy. *Current drug targets*. 2009;10(1):38-41. Epub 2009/01/20.
117. Duffy MJ. The urokinase plasminogen activator system: role in malignancy. *Current pharmaceutical design*. 2004;10(1):39-49. Epub 2004/02/03.
118. Najy AJ, Won JJ, Movilla LS, Kim HR. Differential tumorigenic potential and matriptase activation between PDGF B versus PDGF D in prostate cancer. *Molecular cancer research : MCR*. 2012;10(8):1087-97. Epub 2012/06/13.
119. Li H, Fredriksson L, Li X, Eriksson U. PDGF-D is a potent transforming and angiogenic growth factor. *Oncogene*. 2003;22(10):1501-10. Epub 2003/03/12.
120. Ustach CV, Taube ME, Hurst NJ, Jr., Bhagat S, Bonfil RD, Cher ML, et al. A potential oncogenic activity of platelet-derived growth factor d in prostate cancer progression. *Cancer research*. 2004;64(5):1722-9. Epub 2004/03/05.
121. Wang Z, Kong D, Banerjee S, Li Y, Adsay NV, Abbruzzese J, et al. Down-regulation of platelet-derived growth factor-D inhibits cell growth and angiogenesis through inactivation of Notch-1 and nuclear factor-kappaB signaling. *Cancer research*. 2007;67(23):11377-85. Epub 2007/12/07.
122. Wang Y, Hu C, Dong R, Huang X, Qiu H. Platelet-derived growth factor-D promotes ovarian cancer invasion by regulating matrix metalloproteinases 2 and 9. *Asian Pacific journal of cancer prevention : APJCP*. 2011;12(12):3367-70. Epub 2011/01/01.

123. Cadamuro M, Nardo G, Indraccolo S, Dall'olmo L, Sambado L, Moserle L, et al. Platelet-derived growth factor-D and Rho GTPases regulate recruitment of cancer-associated fibroblasts in cholangiocarcinoma. *Hepatology*. 2013;58(3):1042-53. Epub 2013/03/19.
124. Qin Z, Feng J, Liu Y, Deng LL, Lu C. PDGF-D promotes dermal fibroblast invasion in 3-dimensional extracellular matrix via Snail-mediated MT1-MMP upregulation. *Tumour biology : the journal of the International Society for Oncodevelopmental Biology and Medicine*. 2015. Epub 2015/08/04.
125. Pietras K, Rubin K, Sjoblom T, Buchdunger E, Sjoquist M, Heldin CH, et al. Inhibition of PDGF receptor signaling in tumor stroma enhances antitumor effect of chemotherapy. *Cancer research*. 2002;62(19):5476-84. Epub 2002/10/03.
126. Falcon BL, Pietras K, Chou J, Chen D, Sennino B, Hanahan D, et al. Increased vascular delivery and efficacy of chemotherapy after inhibition of platelet-derived growth factor-B. *The American journal of pathology*. 2011;178(6):2920-30. Epub 2011/06/07.
127. Zhao L, Zhang C, Liao G, Long J. RNAi-mediated inhibition of PDGF-D leads to decreased cell growth, invasion and angiogenesis in the SGC-7901 gastric cancer xenograft model. *Cancer biology & therapy*. 2010;9(1):42-8. Epub 2010/01/21.
128. Liu J, Liao S, Huang Y, Samuel R, Shi T, Naxerova K, et al. PDGF-D improves drug delivery and efficacy via vascular normalization, but promotes lymphatic metastasis by activating CXCR4 in breast cancer. *Clinical cancer research : an official journal of the American Association for Cancer Research*. 2011;17(11):3638-48. Epub 2011/04/05.
129. Kang Y, Massague J. Epithelial-mesenchymal transitions: twist in development and metastasis. *Cell*. 2004;118(3):277-9. Epub 2004/08/06.
130. Kong D, Wang Z, Sarkar SH, Li Y, Banerjee S, Saliganan A, et al. Platelet-derived growth factor-D overexpression contributes to epithelial-mesenchymal transition of PC3 prostate cancer cells. *Stem Cells*. 2008;26(6):1425-35. Epub 2008/04/12.
131. Thiery JP, Acloque H, Huang RY, Nieto MA. Epithelial-mesenchymal transitions in development and disease. *Cell*. 2009;139(5):871-90. Epub 2009/12/01.
132. Wang Z, Ahmad A, Li Y, Kong D, Azmi AS, Banerjee S, et al. Emerging roles of PDGF-D signaling pathway in tumor development and progression. *Biochimica et biophysica acta*. 2010;1806(1):122-30. Epub 2010/05/04.
133. Kong D, Li Y, Wang Z, Banerjee S, Ahmad A, Kim HR, et al. miR-200 regulates PDGF-D-mediated epithelial-mesenchymal transition, adhesion, and invasion of prostate cancer cells. *Stem Cells*. 2009;27(8):1712-21. Epub 2009/06/23.
134. Sethi S, Macoska J, Chen W, Sarkar FH. Molecular signature of epithelial-mesenchymal transition (EMT) in human prostate cancer bone metastasis. *American journal of translational research*. 2010;3(1):90-9. Epub 2010/12/09.
135. Wang R, Li Y, Hou Y, Yang Q, Chen S, Wang X, et al. The PDGF-D/miR-106a/Twist1 pathway orchestrates epithelial-mesenchymal transition in gemcitabine resistance hepatoma cells. *Oncotarget*. 2015;6(9):7000-10. Epub 2015/03/12.
136. Wang Z, Ali S, Banerjee S, Bao B, Li Y, Azmi AS, et al. Activated K-Ras and INK4a/Arf deficiency promote aggressiveness of pancreatic cancer by induction of EMT consistent with cancer stem cell phenotype. *Journal of cellular physiology*. 2013;228(3):556-62. Epub 2012/07/19.
137. Wang Y, Qiu H, Hu W, Li S, Yu J. Over-expression of platelet-derived growth factor-D promotes tumor growth and invasion in endometrial cancer. *International journal of molecular sciences*. 2014;15(3):4780-94. Epub 2014/03/22.
138. Devarajan E, Song YH, Krishnappa S, Alt E. Epithelial-mesenchymal transition in breast cancer lines is mediated through PDGF-D released by tissue-resident stem cells. *International journal of cancer*. 2012;131(5):1023-31. Epub 2011/11/01.
139. Zachary I. Neuropilins: role in signalling, angiogenesis and disease. *Chemical immunology and allergy*. 2014;99:37-70. Epub 2013/11/13.
140. Dhar K, Dhar G, Majumder M, Haque I, Mehta S, Van Veldhuizen PJ, et al. Tumor cell-derived PDGF-B potentiates mouse mesenchymal stem cells-pericytes transition and recruitment through an interaction with NRP-1. *Molecular cancer*. 2010;9:209. Epub 2010/08/07.
141. Evans IM, Yamaji M, Britton G, Pellet-Many C, Lockie C, Zachary IC, et al. Neuropilin-1 signaling through p130Cas tyrosine phosphorylation is essential for growth factor-dependent migration of glioma and endothelial cells. *Molecular and cellular biology*. 2011;31(6):1174-85. Epub 2011/01/20.

142. Cao S, Yaqoob U, Das A, Shergill U, Jagavelu K, Huebert RC, et al. Neuropilin-1 promotes cirrhosis of the rodent and human liver by enhancing PDGF/TGF-beta signaling in hepatic stellate cells. *The Journal of clinical investigation*. 2010;120(7):2379-94. Epub 2010/06/26.
143. Pellet-Many C, Frankel P, Evans IM, Herzog B, Junemann-Ramirez M, Zachary IC. Neuropilin-1 mediates PDGF stimulation of vascular smooth muscle cell migration and signalling via p130Cas. *The Biochemical journal*. 2011;435(3):609-18. Epub 2011/02/11.
144. Pellet-Many C, Mehta V, Fields L, Mahmoud M, Lowe V, Evans I, et al. Neuropilins 1 and 2 mediate neointimal hyperplasia and re-endothelialization following arterial injury. *Cardiovascular research*. 2015;108(2):288-98. Epub 2015/09/28.
145. Pellet-Many C, Frankel P, Jia H, Zachary I. Neuropilins: structure, function and role in disease. *The Biochemical journal*. 2008;411(2):211-26. Epub 2008/03/28.
146. Prud'homme GJ, Glinka Y. Neuropilins are multifunctional coreceptors involved in tumor initiation, growth, metastasis and immunity. *Oncotarget*. 2012;3(9):921-39. Epub 2012/09/06.
147. Gelfand MV, Hagan N, Tata A, Oh WJ, Lacoste B, Kang KT, et al. Neuropilin-1 functions as a VEGFR2 co-receptor to guide developmental angiogenesis independent of ligand binding. *eLife*. 2014;3:e03720. Epub 2014/09/23.
148. Gu C, Limberg BJ, Whitaker GB, Perman B, Leahy DJ, Rosenbaum JS, et al. Characterization of neuropilin-1 structural features that confer binding to semaphorin 3A and vascular endothelial growth factor 165. *The Journal of biological chemistry*. 2002;277(20):18069-76. Epub 2002/03/12.
149. Parker MW, Xu P, Li X, Vander Kooi CW. Structural basis for selective vascular endothelial growth factor-A (VEGF-A) binding to neuropilin-1. *The Journal of biological chemistry*. 2012;287(14):11082-9. Epub 2012/02/10.
150. Herzog B, Pellet-Many C, Britton G, Hartzoulakis B, Zachary IC. VEGF binding to NRP1 is essential for VEGF stimulation of endothelial cell migration, complex formation between NRP1 and VEGFR2, and signaling via FAK Tyr407 phosphorylation. *Molecular biology of the cell*. 2011;22(15):2766-76. Epub 2011/06/10.
151. Wild JR, Staton CA, Chapple K, Corfe BM. Neuropilins: expression and roles in the epithelium. *International journal of experimental pathology*. 2012;93(2):81-103. Epub 2012/03/15.
152. Lampropoulou A, Ruhrberg C. Neuropilin regulation of angiogenesis. *Biochemical Society transactions*. 2014;42(6):1623-8. Epub 2014/11/18.
153. Kitsukawa T, Shimizu M, Sanbo M, Hirata T, Taniguchi M, Bekku Y, et al. Neuropilin-semaphorin III/D-mediated chemorepulsive signals play a crucial role in peripheral nerve projection in mice. *Neuron*. 1997;19(5):995-1005. Epub 1997/12/09.
154. Gerhardt H, Ruhrberg C, Abramsson A, Fujisawa H, Shima D, Betsholtz C. Neuropilin-1 is required for endothelial tip cell guidance in the developing central nervous system. *Developmental dynamics : an official publication of the American Association of Anatomists*. 2004;231(3):503-9. Epub 2004/09/18.
155. Kawasaki T, Kitsukawa T, Bekku Y, Matsuda Y, Sanbo M, Yagi T, et al. A requirement for neuropilin-1 in embryonic vessel formation. *Development*. 1999;126(21):4895-902. Epub 1999/10/16.
156. Kitsukawa T, Shimono A, Kawakami A, Kondoh H, Fujisawa H. Overexpression of a membrane protein, neuropilin, in chimeric mice causes anomalies in the cardiovascular system, nervous system and limbs. *Development*. 1995;121(12):4309-18. Epub 1995/12/01.
157. Lanahan A, Zhang X, Fantin A, Zhuang Z, Rivera-Molina F, Speichinger K, et al. The neuropilin 1 cytoplasmic domain is required for VEGF-A-dependent arteriogenesis. *Developmental cell*. 2013;25(2):156-68. Epub 2013/05/04.
158. Fantin A, Schwarz Q, Davidson K, Normando EM, Denti L, Ruhrberg C. The cytoplasmic domain of neuropilin 1 is dispensable for angiogenesis, but promotes the spatial separation of retinal arteries and veins. *Development*. 2011;138(19):4185-91. Epub 2011/08/20.
159. Kofler N, Simons M. The expanding role of neuropilin: regulation of transforming growth factor-beta and platelet-derived growth factor signaling in the vasculature. *Current opinion in hematology*. 2016. Epub 2016/02/06.
160. Aspalter IM, Gordon E, Dubrac A, Ragab A, Narloch J, Vizan P, et al. Alk1 and Alk5 inhibition by Nrp1 controls vascular sprouting downstream of Notch. *Nature communications*. 2015;6:7264. Epub 2015/06/18.
161. Glinka Y, Prud'homme GJ. Neuropilin-1 is a receptor for transforming growth factor beta-1, activates its latent form, and promotes regulatory T cell activity. *Journal of leukocyte biology*. 2008;84(1):302-10. Epub 2008/04/26.

162. Glinka Y, Stoilova S, Mohammed N, Prud'homme GJ. Neuropilin-1 exerts co-receptor function for TGF-beta-1 on the membrane of cancer cells and enhances responses to both latent and active TGF-beta. *Carcinogenesis*. 2011;32(4):613-21. Epub 2010/12/28.
163. Di Guglielmo GM, Le Roy C, Goodfellow AF, Wrana JL. Distinct endocytic pathways regulate TGF-beta receptor signalling and turnover. *Nature cell biology*. 2003;5(5):410-21. Epub 2003/04/30.
164. Valdembri D, Caswell PT, Anderson KI, Schwarz JP, Konig I, Astanina E, et al. Neuropilin-1/GIPC1 signaling regulates alpha5beta1 integrin traffic and function in endothelial cells. *PLoS biology*. 2009;7(1):e25. Epub 2009/01/30.
165. Koch S, van Meeteren LA, Morin E, Testini C, Westrom S, Bjorkelund H, et al. NRP1 presented in trans to the endothelium arrests VEGFR2 endocytosis, preventing angiogenic signaling and tumor initiation. *Developmental cell*. 2014;28(6):633-46. Epub 2014/03/25.
166. Frankel P, Pellet-Many C, Lehtolainen P, D'Abaco GM, Tickner ML, Cheng L, et al. Chondroitin sulphate-modified neuropilin 1 is expressed in human tumour cells and modulates 3D invasion in the U87MG human glioblastoma cell line through a p130Cas-mediated pathway. *EMBO reports*. 2008;9(10):983-9. Epub 2008/08/16.
167. Shintani Y, Takashima S, Asano Y, Kato H, Liao Y, Yamazaki S, et al. Glycosaminoglycan modification of neuropilin-1 modulates VEGFR2 signaling. *The EMBO journal*. 2006;25(13):3045-55. Epub 2006/06/10.
168. Xian X, Hakansson J, Stahlberg A, Lindblom P, Betsholtz C, Gerhardt H, et al. Pericytes limit tumor cell metastasis. *The Journal of clinical investigation*. 2006;116(3):642-51. Epub 2006/02/14.
169. Bergers G, Song S, Meyer-Morse N, Bergsland E, Hanahan D. Benefits of targeting both pericytes and endothelial cells in the tumor vasculature with kinase inhibitors. *The Journal of clinical investigation*. 2003;111(9):1287-95. Epub 2003/05/03.
170. Ballmer-Hofer K, Andersson AE, Ratcliffe LE, Berger P. Neuropilin-1 promotes VEGFR-2 trafficking through Rab11 vesicles thereby specifying signal output. *Blood*. 2011;118(3):816-26. Epub 2011/05/19.